### bs-23703R

## [ Primary Antibody ]

# BIOSS ANTIBODIES

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

IHC-F (1:100-500)

**IF** (1:100-500)

Reactivity: Mouse (predicted: Human,

Rat, Rabbit, Pig, Sheep, Cow, Dog, Horse)

Applications: IHC-P (1:100-500)

Predicted 127 kDa

Subcellular Location: Cell membrane

# **ERG/KCNH2** Rabbit pAb

- DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**GenelD:** 3757 **SWISS:** Q12809

Target: ERG/KCNH2

**Immunogen:** KLH conjugated synthetic peptide derived from human

ERG/KCNH2: 751-850/1159. < Cytoplasmic >

**Purification:** affinity purified by Protein A

Concentration: 1mg/ml

**Storage:** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.


**Background:** The potassium voltage gated channel, subfamily H (eag related),

member 2 (KCNH2) gene encodes a voltage-gated potassium channel which has an important role in cardiac action potential repolarization in the mammalian heart. Mutations in KCNH2 have been shown to cause chromosome 7-linked congenital long QT syndrome, a disorder associated with delayed cardiac repolarization, prolonged electrocardiographic QT intervals, and

the development of ventricular arrhythmias. KCNH2 channels are an important target for many drugs, and have emerged as a significant type of cardiac ion channel. Highly expressed in heart

and brain.

VALIDATION IMAGES



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (ERG/KCNH2) Polyclonal Antibody, Unconjugated (bs-23703R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

#### SELECTED CITATIONS —

• [IF=3.499] Zhan C et al. Rotenone and 3-bromopyruvate toxicity impacts electrical and structural cardiac remodeling in rats. Toxicol Lett. 2019 Oct 1. pii: S0378-4274(19)30295-4. IHC; Rat. 31585160