

bs-11478R**[Primary Antibody]****SEMA4F Rabbit pAb**

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —

Host: Rabbit	Isotype: IgG	Applications: WB (1:500-2000) IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) ICC/IF (1:100-500) ELISA (1:5000-10000) Reactivity: (predicted: Human, Mouse, Rat, Cow, Dog, Horse) Predicted MW.: 80 kDa Subcellular Location: Cell membrane
Clonality: Polyclonal		
GeneID: 10505	SWISS: O95754	
Target: SEMA4F		
Immunogen: KLH conjugated synthetic peptide derived from human SEMA4F: 239-288/770. < Extracellular >		
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: Semaphorins are a family of cell surface and secreted proteins that are conserved from insects to humans. Members of this family of proteins are approximately 750 amino acids in length (including signal sequences) and are defined by a conserved extracellular “semaphorin” domain of approximately 500 amino acids containing 14-16 cysteines, blocks of conserved sequences and no obvious repeats. Secreted and cell-bound semaphorins chemically attract and repel the growth of neural axons, guiding the development of intricate networks of neural tissue. SEMA4F (semaphorin-4F), also known as SEMAM, SEMAW or PRO2353, is a 770 amino acid member of the semaphorin family. Localized to the membrane, SEMA4F is a single-pass type I protein that is involved in growth cone collapse of retinal ganglion-cell axons. SEMA4F is highly expressed in postnatal brain and lung and contains one immunoglobulin-like (Ig-like) domain, one PSI domain and one semaphorin domain. Two isoforms exist due to alternative splicing events.		