

**bs-20273R****[ Primary Antibody ]****ADAM2 Rabbit pAb****BioSS**  
**ANTIBODIES**

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

**— DATASHEET —**

<b>Host:</b> Rabbit <b>Clonality:</b> Polyclonal <b>GeneID:</b> 2515 <b>Target:</b> ADAM2 <b>Immunogen:</b> KLH conjugated synthetic peptide derived from human ADAM2: 451-550/735. < Extracellular > <b>Purification:</b> affinity purified by Protein A <b>Concentration:</b> 1mg/ml <b>Storage:</b> Preservative: 0.02% Proclin300, Constituents: 1% BSA, 0.01M PBS, pH7.4. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. <b>Background:</b> ADAM-2, also known as PH-30 and Fertilin-2 was first described as a sperm-egg fusion protein from guinea pig. A member of the metalloproteinase family containing disintegrin-like domains (ADAMs), the function of ADAM-2 is still poorly understood. Unlike ADAM-1, ADAM-2 does not contain the canonical HEXXHxxxxH zinc metalloproteinase motif, and is not thought to be proteolytically active. Like the other ADAMs, ADAM-2 domain structure consists of a signal sequence followed by a propeptide domain, a metalloproteinase domain, a disintegrin domain cysteine-rich domain, an EGF-like domain, a transmembrane domain, and a cytoplasmic domain. Three isoforms of ADAM-2 are reported to date, which differ in the beginning of the metalloproteinase domain and cysteine-rich domain. The sequences are coded from intronless genes. The longest ADAM-2 message encodes a protein of 735 amino acids, with a predicted mass of 82.5 kDa and a pI of 5.73. The 716 amino acid form of ADAM-2 has a deletion in the start of the metalloproteinase domain, relative to the longer form, and has a predicted mass of 80.2 kDa and pI of 5.75. The shorter form of 579 amino acids shared the deletion in the MP domain, and also has a deletion in the cysteine-rich domain. Localized on the surface of sperm, the ADAM-2 isoforms are thought to form a heterodimer with ADAM-1 (fertilin-a), and facilitate sperm-egg fusion, although there is some controversy about the precise actions the proteins play. Integrin $\alpha 6$ on the egg surface is thought to dock with a QDECD motif in the disintegrin domain of ADAM-2, and there is some speculation that ADAM-1/ADAM-2 heterodimer initiated ADAM-3 production on the cell surface.	<b>Isotype:</b> IgG <b>SWISS:</b> Q99965 <b>Applications:</b> WB (1:500-2000) <b>Reactivity:</b> Human  <b>Predicted MW.:</b> 63/80 kDa <b>Subcellular Location:</b> Cell membrane
---	--

**— VALIDATION IMAGES —**

Sample: k562 (Human) Cell Lysate at 40 ug  
 Primary: Anti-RNF14(bs-20273R) at 1/300 dilution  
 Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 63 kD  
 Observed band size: 63 kD