bs-20503R

[Primary Antibody]

PFKP Rabbit pAb



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

- DATASHEET		400-901-9800
Host: Rabbit	Isotype: IgG	Applications: IHC-P (1:100-500) IHC-F (1:100-500)
Clonality: Polyclonal GenelD: 5214	SWISS: Q01813	IF (1:100-500) ICC/IF (1:100-500) ELISA (1:5000-10000)
Target: PFKP Immunogen: KLH conjugated synthetic peptide derived from human PFKP: 231-330/784.		Reactivity: (predicted: Human, Mouse, Rat)
Purification: affinity purified by P	rotein A	
Concentration: 1mg/ml		Predicted 86 kDa
Storage: Preservative: 0.02% Proclin300, Constituents: 1% BSA, 0.01M PBS, pH7.4. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		Subcellular Location: Cytoplasm
phosphotransferase conversion of fructo and is a key regulato maps to chromosom the muscle (PFKM; M isoforms of phospho 12q13 and 21q22, re determined that full	e (PFK) (ATP:D-fructose-6-phosphate-1- , EC 2.7.1.11). PFK catalyzes the irreversible se-6-phosphate to fructose-1,6-bisphosphate ry enzyme in glycolysis. The PFKP gene, whic ie 10p, is also expressed in fibroblasts. See al IIM 610681) and liver (PFKL; MIM 171860) fructokinase, which map to chromosomes spectively. Vora (1981) [PubMed 6451249] tetrameric phophofructokinase enzyme ts can be composed of subunits P4, P3L, and	ch so