

bsm-51385M**[Primary Antibody]****EIF2AK3/PERK Mouse mAb****BioSS**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —

Host: Mouse Clonality: Monoclonal GeneID: 9451 Target: EIF2AK3/PERK Purification: affinity purified by Protein G Concentration: 1mg/ml Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. Background: The protein encoded by this gene phosphorylates the alpha subunit of eukaryotic translation-initiation factor 2 (EIF2), leading to its inactivation, and thus to a rapid reduction of translational initiation and repression of global protein synthesis. It is a type I membrane protein located in the endoplasmic reticulum (ER), where it is induced by ER stress caused by malformed proteins. Mutations in this gene are associated with Wolcott-Rallison syndrome. [provided by RefSeq, Jan 2010].	Isotype: IgG1 CloneNo.: 3C3 SWISS: Q9NZJ5	Applications: WB (1:500-2000) ELISA (1:5000-10000) Reactivity: Mouse, Rat (predicted: Human) Predicted MW.: 122 kDa Subcellular Location: Endoplasmic reticulum, Membrane
--	--	--

— VALIDATION IMAGES —

25 ug total protein per lane of various lysates (see on figure) probed with EIF2AK3/PERK monoclonal antibody, unconjugated (bsm-51385M) at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.

— SELECTED CITATIONS —

- **[IF=5.793]** Chunyue Wang. et al. Neuroprotective effects of verbascoside against Alzheimer' s disease via the relief of endoplasmic reticulum stress in Aβ-exposed U251 cells and APP/PS1 mice. J Neuroinflamm. 2020 Dec;17(1):1-16 WB ;Human. 33070776