## bs-6347R

## [ Primary Antibody ]

## Zinc finger protein 754 Rabbit pAb



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

- DATASHEET		400-901-9800
Host: Rabbit	Isotype: IgG	Applications: WB (1:500-2000)
Clonality: Polyclonal		Reactivity: Human, Mouse
GenelD: 132625		(predicted: Rat, Rabbit, Pig,
Target: Zinc finger protein 754		Sheep, Cow, Horse)
Immunogen: KLH conjugated synthetic peptide derived from human Rex1/Zinc finger protein 754: 191-290/310.		Predicted MW.: <sup>35 kDa</sup>
Purification: affinity purified by Pr	otein A	Subcellular
Concentration: 1mg/ml		Subcellular Location: <sup>Nucleus</sup>
Glycerol.	th 1% BSA, 0.02% Proclin300 and 50% e at -20°C for one year. Avoid repeated	
involved in transcript Expressed in kidney, cells, bronchial and s level). Expressed in n lines (at protein level epidermal keratinocy small airway lung epi	wal property of ES cells (By similarity). I tional regulation. Tissue specificity: epidermal keratinocytes, prostate epit mall airway lung epithelial cells (at pro halignant kidney and several carcinom ). Expressed in embryonic stem cells, k ytes, prostate epithelial cells, bronchia ithelial cells. Expressed in embryonal nas, malignant kidney and several carc	thelial otein a cell kidney, l and

## - VALIDATION IMAGES -



Sample: HL60(Human) Cell Lysate at 30 ug Primary: Anti- Zinc finger protein 754 (bs-6347R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 35 kD Observed band size: 35 kD Sample: NIH/3T3(Mouse) Cell Lysate at 30 ug Primary: Anti- Zinc finger protein 754 (bs-6347R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 35 kD Observed band size: 35 kD