

---

## IRX1 Rabbit pAb

Catalog Number: bs-11857R

Target Protein: IRX1

Concentration: 1mg/ml

Form: Liquid

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: IHC-P (1:100-500), IHC-F (1:100-500), IF (1:100-500)

Reactivity: Mouse, Rat (predicted:Human, Rabbit, Cow, Chicken, Dog, Horse)

Predicted MW: 50 kDa

Entrez Gene: 79192

Swiss Prot: P78414

Source: KLH conjugated synthetic peptide derived from human IRX1: 166-240/480.

Purification: affinity purified by Protein A


Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.


**Background:** The Iroquois homeobox gene family of transcription factors regulate aspects of embryonic development including anterior/posterior and dorsal/ventral axis patterning in the central nervous system. The Iroquois family are clustered on two loci, IRXA and IRXB, which map to chromosomes 8 and 13 in mice. The IRXA group includes *Irx1*, *Irx2* and *Irx4*; the IRXB group is comprises *Irx3*, *Irx5* and *Irx6*. *Irx1* and *Irx2* are both widely expressed during development in the lung epithelium and also in the ventricular septum. *Irx1* and *Irx2* also play a role in digit formation (E11.5–E14.5). The *Irx* gene family members are each expressed in a distinct pattern during mouse heart development. Specifically, *Irx1* and *Irx2* are expressed in the ventricular septum and *Irx3* is expressed in the ventricular trabeculated myocardium. In addition, *Irx4* is expressed in the linear heart tube and the AV canal; *Irx5* is expressed in the endocardium lining the ventricular and atrial myocardium. Furthermore, the *IRX4* gene may modulate cardiac development and function. Although the heart of *Irx4*(-) mice appears to develop normally, adult *Irx4*(-) mice exhibit cardiomyopathy, including cardiac hypertrophy and decreased contractility.

## VALIDATION IMAGES


---



Tissue/cell: rat brain tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Incubation: Anti-IRX1 Polyclonal Antibody, Unconjugated (bs-11857R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody (SP-0023) and DAB (C-0010) staining



Paraformaldehyde-fixed, paraffin embedded (mouse intestine); Antigen retrieval by boiling in sodium citrate buffer (pH 6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (IRX) Polyclonal Antibody, Unconjugated (bs-11857R) at 1:200 overnight at 4°C, followed by operating according to SP Kit (Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH 6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (IRX) Polyclonal Antibody, Unconjugated (bs-11857R) at 1:200 overnight at 4°C, followed by operating according to SP Kit (Rabbit) (sp-0023) instructions and DAB staining.