

bs-11858R**[Primary Antibody]****ITM2B Rabbit pAb****BioSS**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —

Host: Rabbit	Isotype: IgG	Applications: IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500)
Clonality: Polyclonal		
GeneID: 9445	SWISS: Q9Y287	
Target: ITM2B		Reactivity: Rat (predicted: Human, Mouse)
Immunogen: KLH conjugated synthetic peptide derived from human ITM2B: 10-90/266.		
Purification: affinity purified by Protein A		Predicted MW.: 30 kDa
Concentration: 1mg/ml		Subcellular Location: Cell membrane ,Cytoplasm
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: The type II integral membrane (ITM2) protein family consists of three members: ITM2A (also designated E25), ITM2B and ITM2C. ITM2A expression is high in osteogenic and lymphoid tissues, while both ITM2B and ITM2C are expressed in brain. ITM2B is a 266 amino acid protein that contains a potential N-glycosylation site, a potential single transmembrane-spanning domain between amino acids 52 and 74 and an extracellular C-terminal domain. Mutations in the ITM2B gene can lead to familial British dementia (FBD), and autosomal dominant disease with an onset around the fifth decade of life that is characterized by progressive dementia, spasticity and cerebellar ataxia. Familial Danish dementia (FDD), also designated hereditary ophthalmic-oto-encephalica, is also associated with mutations in the ITM2B gene. FDD is an autosomal dominant disorder characterized by cataracts, deafness, progressive ataxia and dementia.		

— VALIDATION IMAGES —

Paraformaldehyde-fixed, paraffin embedded (rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (ITM2B) Polyclonal Antibody, Unconjugated (bs-11858R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.