

bs-7721R

[Primary Antibody]

CAPG2 Rabbit pAb

Bioss
ANTIBODIES

www.bioss.com.cn

sales@bioss.com.cn


techsupport@bioss.com.cn

400-901-9800


DATASHEET

Host: Rabbit	Isotype: IgG	Applications: WB (1:500-2000) IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) Flow-Cyt (2ug/Test)
Clonality: Polyclonal		
GeneID: 54892	SWISS: Q86XI2	
Target: CAPG2		
Immunogen: KLH conjugated synthetic peptide derived from human CAPG2/NCAPG2: 451-550/1143.		
Purification: affinity purified by Protein A		Reactivity: Human, Mouse, Rat
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		Predicted MW.: 131 kDa
Background: NCAPG2 is 1 of 3 non-SMC subunits that define condensin II. Condensin complexes I and II play essential roles in mitotic chromosome assembly and segregation. Both condensins contain 2 invariant structural maintenance of chromosome (SMC) subunits, SMC2 and SMC4, but they contain different sets of non-SMC subunits. NCAPG2 is 1 of 3 non-SMC subunits that defines condensin II.		Subcellular Location: Nucleus


VALIDATION IMAGES




Sample: Lane 1: Mouse Thymus tissue lysates
Lane 2: Rat Testis tissue lysates Lane 3: Human A431 cell lysates Lane 4: Human MCF-7 cell lysates Lane 5: Human HeLa cell lysates Lane 6: Human U251 cell lysates Lane 7: Human Jurkat cell lysates Primary: Anti-CAPG2 (bs-7721R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 131 kDa Observed band size: 131 kDa



Sample: 293T(Human) Cell Lysate at 30 ug
Primary: Anti-CAPG2 (bs-7721R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 131 kD Observed band size: 131 kD



Paraformaldehyde-fixed, paraffin embedded (human tonsil); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Incubation with (CAPG2) Polyclonal Antibody, Unconjugated (bs-7721R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Blank control: Mouse spleen. Primary Antibody (green line): Rabbit Anti-CAPG2 antibody (bs-7721R) Dilution: 2µg / 10⁶ cells; Isotype Control Antibody (orange line): Rabbit IgG . Secondary Antibody : Goat anti-rabbit IgG-AF647 Dilution: 1µg / test. Protocol The cells were fixed with 4% PFA (10min at room temperature) and then permeabilized with 90% ice-cold methanol

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

for 20 min at 20°C. The cells were then incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at room temperature .Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.

— SELECTED CITATIONS —

- **[IF=64.8]** El Yakoubi Warif. et al. Condensin dysfunction is a reproductive isolating barrier in mice. NATURE. 2023 Nov;;1-8 IF,WB ;Mouse. 37914934
- **[IF=8.44]** Wang Wei. et al. Circular RNA circ0001955 promotes cervical cancer tumorigenesis and metastasis via the miR-188-3p/NCAPG2 axis. J TRANSL MED. 2023 Dec;21(1):1-18 IHC,WB ;Human,Mouse. 37248471