

bsm-52146R**[Primary Antibody]****BioSS**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

phospho-Cyclin E1 (Thr77) Recombinant Rabbit mAb**DATASHEET**


Host: Rabbit	Isotype: IgG	Applications: WB (1:200-500) IHC-P (1:100-500) IHC-F (1:50-200) IF (1:50-200) ICC/IF (1:50-200)
Clonality: Recombinant	CloneNo.: 4G10	
GeneID: 898	SWISS: P24864	
Target: phospho-Cyclin E1 (Thr77)		
Immunogen: KLH conjugated Synthesised phosphopeptide derived from human Cyclin E around the phosphorylation site of Thr77: IP(p-T)PD.		
Purification: affinity purified by Protein A		Reactivity: Human, Mouse, Rat
Concentration: 1mg/ml		Predicted MW.: 47 kDa
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: Cyclin E is a regulatory subunit of Cdk2 and controls G1 / S transition during the mammalian cell cycle. Multiple isoforms of Cyclin E are only expressed in tumors but not in normal tissue, suggesting a post transcriptional regulation of Cyclin E. In vitro analyses indicated that these truncated variant isoforms of Cyclin E are able to phosphorylate histone H1. Alterations in the Cyclin E protein have been implicated as indicators of worse prognosis in various cancers.		Subcellular Location: Nucleus

VALIDATION IMAGES


Immunohistochemical analysis of paraffin-embedded human testis tissue using anti-Phospho-Cyclin E1(T77) antibody. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0) for 20 minutes. The tissues were blocked in 1% BSA for 30 minutes at room temperature, washed with ddH2O and PBS, and then probed with the primary antibody (bsm-52146R, 1/400) for 30 minutes at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.



Immunohistochemical analysis of paraffin-embedded human tonsil tissue using anti-Phospho-Cyclin E1(T77) antibody. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0) for 20 minutes. The tissues were blocked in 5% BSA for 30 minutes at room temperature, washed with ddH2O and PBS, and then probed with the primary antibody (bsm-52146R, 1/50) for 30 minutes at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.



ICC staining of Phospho-Cyclin E1(T77) in HepG2 cells (green). Formalin fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 10 minutes at room temperature and blocked with 1% Blocker BSA for 15 minutes at room temperature. Cells were probed with the primary antibody (bsm-52146R, 1/50) for 1 hour at room temperature, washed with PBS. Alexa Fluor®488 Goat anti-Rabbit IgG was used as the secondary antibody at 1/1,000 dilution. The nuclear counter stain is DAPI (blue).



ICC staining of Phospho-Cyclin E1(T77) in HeLa cells (green). Formalin fixed cells were permeabilized with 0.1% Triton X-100 in TBS for

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

10 minutes at room temperature and blocked with 1% Blocker BSA for 15 minutes at room temperature. Cells were probed with the primary antibody (bsm-52146R, 1/50) for 1 hour at room temperature, washed with PBS. Alexa Fluor®488 Goat anti-Rabbit IgG was used as the secondary antibody at 1/1,000 dilution. The nuclear counter stain is DAPI (blue).