bs-1819R

[Primary Antibody]

Bioss

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

HDAC3 Rabbit pAb

- DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

GenelD: 8841 **SWISS:** 015379

Target: HDAC3

Immunogen: KLH conjugated synthetic peptide derived from human HDAC3:

351-428/428.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: Histones play a critical role in transcriptional regulation, cell cycle

progression, and developmental events. Histone

acetylation/deacetylation alters chromosome structure and affects transcription factor access to DNA. The protein encoded by this gene belongs to the histone deacetylase/acuc/apha family. It has histone deacetylase activity and represses transcription when tethered to a promoter. It may participate in the regulation of transcription through its binding with the zinc-finger transcription factor YY1. This protein can also down-regulate p53 function and thus modulate cell growth and apoptosis. This gene is regarded as a potential tumor suppressor gene. [provided by RefSeq, Jul 2008]

Applications: WB (1:500-2000)

IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) ICC/IF (1:100-500)


Reactivity: Human, Mouse, Rat

(predicted: Pig, Cow, Chicken, Horse)

Predicted MW.: 47 kDa

Subcellular Location: Cell membrane ,Nucleus


VALIDATION IMAGES



Sample: Uterus (Mouse) Lysate at 40 ug Ovary (Mouse) Lysate at 40 ug Primary: Anti-HDAC3 (bs-1819R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 47 kD Observed band size: 50 kD




Sample: Lane 1: Mouse Cerebrum tissue lysates Lane 2: Rat Cerebrum tissue lysates Primary: Anti-HDAC3/HD3 (bs-1819R) at 1/500 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 47 kDa Observed band size: 47 kDa




Paraformaldehyde-fixed, paraffin embedded (human gastric carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (HDAC3) Polyclonal Antibody, Unconjugated (bs-1819R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructionsand DAB staining.



Tissue/cell: rat brain tissue; 4% Paraformaldehyde-fixed and paraffinembedded; Antigen retrieval: citrate buffer (



Tissue/cell: rat brain tissue;4% Paraformaldehyde-fixed and paraffinembedded; Antigen retrieval: citrate buffer (



issue/cell: rat brain tissue;4% Paraformaldehyde-fixed and paraffinembedded; Antigen retrieval: citrate buffer (

0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min; Incubation:
Anti-HDAC3/HD3 Polyclonal Antibody,
Unconjugated(bs-1819R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining

0.01M, pH 6.0), Boiling bathing for 15min;
Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min; Incubation: Anti-HDAC3/HD3
Polyclonal Antibody, Unconjugated(bs-1819R)
1:200, overnight at 4°C; The secondary antibody was Goat Anti-Rabbit IgG, Cy3
conjugated(bs-0295G-Cy3, red)used at 1:200
dilution for 40 minutes at 37°C.

0.01M, pH 6.0), Boiling bathing for 15min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min; Incubation: Anti-HDAC3/HD3 Polyclonal Antibody, Unconjugated(bs-1819R) 1:200, overnight at 4°C; The secondary antibody was Goat Anti-Rabbit IgG, Cy3 conjugated(bs-0295G-Cy3)used at 1:200 dilution for 40 minutes at 37°C. DAPI(5ug/ml,blue,C-0033) was used to stain the cell nuclei

- SELECTED CITATIONS -

- [IF=5] Guifeng Zhang. et al. Sea Cucumber Egg Oligopeptides Ameliorate Cognitive Impairments and Pathology of Alzheimer's Disease Through Regulating HDAC3 and BDNF/NT3 via the Microbiota–Gut–Brain Axis. NUTRIENTS. 2025 Jan;17(14):2312 WB; 40732937
- [IF=4] Dan Pu. et al. Combined supplementation of short-chain fatty acids reduces hyperphosphorylation of Tau at T181,T231 and S396 sites and improves cognitive impairment in a chemically induced AD mouse model via regulation of HDAC and Keap1. NEUROCHEM INT. 2025 Oct;189:106034 WB; Mouse. 40812734