bs-2891R

[Primary Antibody]

BIOSS ANTIBODIES

HDAC8 Rabbit pAb

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

- DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

GenelD: 55869 **SWISS:** Q9BY41

Target: HDAC8

Immunogen: KLH conjugated synthetic peptide derived from human HDAC8:

151-250/377.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: Histones play a critical role in transcriptional regulation, cell cycle

progression, and developmental events. Histone

acetylation/deacetylation alters chromosome structure and affects transcription factor access to DNA. The protein encoded by this gene belongs to class I of the histone deacetylase family. It catalyzes the deacetylation of lysine residues in the histone N-terminal tails and represses transcription in large multiprotein complexes with transcriptional co-repressors. Multiple transcript variants encoding different isoforms have been found for this gene.

[provided by RefSeq, Oct 2009].

Applications: IHC-P (1:100-500)

IHC-F (1:100-500) IF (1:100-500)


Reactivity: Mouse (predicted: Human,

Rat, Rabbit, Pig, Cow, Chicken, Horse)

Predicted MW.: 42 kDa

Subcellular Location: Cytoplasm ,Nucleus

VALIDATION IMAGES



Tissue/cell: mouse pancreas tissue; 4%
Paraformaldehyde-fixed and paraffinembedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min; Incubation: Anti-HDAC8 Polyclonal Antibody, Unconjugated(bs-2891R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining