bs-3489R

[Primary Antibody]

phospho-Tau (Ser422) Rabbit pAb



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

DATASHEET —

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

GenelD: 4137 **SWISS:** P10636

Target: Tau (Ser422)

Immunogen: KLH conjugated Synthesised phosphopeptide derived from human

Tau around the phosphorylation site of Ser422: VD(p-S)PQ.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: Tau proteins are important Promotes microtubule assembly and stability, and might be involved in the establishment and maintenance of $\bar{\text{n}}\text{euronal}$ polarity. The C-terminus binds axonal microtubules while the N-terminus binds neural plasma membrane components, suggesting that tau functions as a linker protein between both. Axonal polarity is predetermined by tau localization (in the neuronal cell) in the domain of the cell body defined by the centrosome. The short isoforms allow plasticity of the cytoskeleton whereas the longer isoforms may preferentially play a role in its stabilization. Tau proteins subcellular located in the axons of neurons, in the cytoso I and in association with plasma membrane components. It expressed in neurons. PNS-tau is expressed in the peripheral nervous system while the others are expressed in the central nervous system.

Applications: WB (1:500-2000)

IHC-P (1:100-500) IHC-F (1:100-500) **IF** (1:100-500)

Reactivity: Mouse, Rat


(predicted: Human, Rabbit,

Cow, Dog)


Predicted 52/79 kDa

Subcellular Location: Cell membrane ,Cytoplasm

VALIDATION IMAGES



Sample: Cerebellum (Mouse) Lysate at 40 ug Cerebrum (Rat) Lysate at 40 ug Lung (Mouse) Lysate at 40 ug Kidney (Mouse) Lysate at 40 ug Primary: Anti- phospho-Tau (Ser422) (bs-3489R) at 1/1000 dilution Secondary: IRDve800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 83 kD Observed band size: 52 kD



Paraformaldehyde-fixed, paraffin embedded (Rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes: Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (phospho-Tau (Ser422)) Polyclonal Antibody, Unconjugated (bs-3489R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

— SELECTED CITATIONS —

• [IF=15.8] Adriena Jedličková. et al. Inhaled Lead Nanoparticles Enter the Brain through the Olfactory Pathway and Induce Neurodegenerative Changes Resembling Tauopathies..ACS Nano.2025 Mar 25. WB, IHC; Mouse, zebrafish. 40130682

- [IF=6.2] Bo Zhang. et al. Arsenic exposure activates microglia, inducing neuroinflammation and promoting the occurrence and development of Alzheimer's disease-like neurodegeneration in mice. ECOTOX ENVIRON SAFE. 2025

 Jun;297:118251 IHC,IF; Mouse. 40294499
- [IF=4] Wang Xiaoping. et al. S1PR2 Regulates Autophagy Through the AKT/mTOR Pathway to Promote Pathological Damage in Alzheimer's Disease. J ALZHEIMERS DIS. 2023 Nov;Preprint(Preprint):1-16 WB;Human. 38007654