

bs-3641R**[Primary Antibody]****BioSS**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

SUR1 Rabbit pAb**— DATASHEET —**

Host: Rabbit Clonality: Polyclonal GeneID: 6833 Target: SUR1 Immunogen: KLH conjugated synthetic peptide derived from human SUR1: 301-400/1581. < Extracellular > Purification: affinity purified by Protein A Concentration: 1mg/ml Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. Background: SUR1 is a member of ATP binding cassette super family. SUR receptor confers the sensitivity of Kir6.2 to ATP/ADP sensitivity and to pharmacological agents such as sulfonylurea and diazoxide that close or open the KATP channels. The persistent hyperinsulinemic hypoglycemia in infancy (PHHI) is familial disorder due to defect in negative feed back in response to low glucose levels. SUR1 was mapped on chromosome 11p14-15.1, the same location where the gene for PHHI is located. It has been shown that the expression of Kir6.2 and SUR1 are regulated by glucose levels and the actives of glucagon like peptide receptor 1. Abnormal insulin secretion in PHHI appears to be caused by mutations in the SUR gene.	Isotype: IgG SWISS: Q09428 Applications: WB (1:500-2000) Reactivity: Human, Mouse, Rat (predicted: Rabbit, Cow, Horse) Predicted MW.: 175 kDa Subcellular Location: Cell membrane
--	---

— VALIDATION IMAGES —

Sample: Hippocampus (Mouse) Lysate at 40 ug
Primary: Anti-SUR1 (bs-3641R) at 1/1000 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 175 kD
Observed band size: 175 kD



Sample: Lane 1: Cerebrum (Mouse) Lysate at 40 ug
Lane 2: Pancreas (Mouse) Lysate at 40 ug
Lane 3: Stomach (Mouse) Lysate at 40 ug
Lane 4: Cerebrum (Rat) Lysate at 40 ug
Lane 5: SH-SY5Y (Human) Cell Lysate at 30 ug
Primary: Anti-SUR1 (bs-22836R) at 1/1000 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution
Predicted band size: 177 kD
Observed band size: 180 kD

— SELECTED CITATIONS —

- **[IF=0.91]** Horii K et al. ATP-dependent potassium channels contribute to motor regulation of esophageal striated muscle in rats. J Vet Med Sci. 2019 Jul 9. IHC ;Rat. 31292350