

bsm-60597R**[Primary Antibody]****BioSS**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

TG Recombinant Rabbit mAb**DATASHEET**


Host: Rabbit	Isotype: IgG	Applications: WB (1:500-2000) IHC-P (1:50-200) IHC-F (1:50-200) IF (1:50-200) Reactivity: Human, Mouse, Rat Predicted MW.: 660 kDa Subcellular Location: Secreted
Clonality: Recombinant	CloneNo.: A7B9	
GeneID: 7038	SWISS: P01266	
Target: TG		
Immunogen: A synthesized peptide derived from human Thyroglobulin: 1001-1023/2768.		
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: Thyroglobulin is the glycoprotein precursor of the iodinated thyroid hormones thyroxine (T4) and triiodothyronine (T3). Thyroglobulin is obtained from the thyroid gland and exhibits the general properties of the globulins. The human thyroglobulin (hTG) is a high molecular weight glycoprotein (605 kDa) found in the thyroid follicular cells. It plays a central role in the uptake, incorporation, and regulated biosynthesis of thyroid hormones, T4 and T3. Thyroid disorders are, in large part, due to autoimmune origin, and anti thyroglobulin autoantibodies were the first factor to be discovered. Anti hTG is found in all thyroid autoimmune diseases (Hashimoto's thyroiditis, Graves' diseases), with the highest level observed in Hashimoto's thyroiditis. Anti hTG is also characteristic of thyroid cancer, and its determination can be used for the follow up of cancer patients.		

VALIDATION IMAGES

25 ug total protein per lane of various lysates (see on figure) probed with TG monoclonal antibody, unconjugated (bsm-60597R) at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.



Paraformaldehyde-fixed, paraffin embedded Human Thyroid Cancer; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with TG Monoclonal Antibody, Unconjugated(bsm-60597R) at 1:200 overnight at 4°C, followed by conjugation to the bs-0295G-HRP and DAB (C-0010) staining.



Paraformaldehyde-fixed, paraffin embedded Rat Thyroid Gland; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with TG Monoclonal Antibody, Unconjugated(bsm-60597R) at 1:200 overnight at 4°C, followed by conjugation to the bs-0295G-HRP and DAB (C-0010) staining.