

**bs-3046R****[ Primary Antibody ]****Phospho-Ack1 (Tyr326) Rabbit pAb**

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

**— DATASHEET —**

<p><b>Host:</b> Rabbit</p> <p><b>Clonality:</b> Polyclonal</p> <p><b>GeneID:</b> 10188</p> <p><b>Target:</b> Phospho-Ack1 (Tyr326)</p> <p><b>Immunogen:</b> KLH conjugated synthetic peptide derived from human Ack1: FT(p-Y)GQ.</p> <p><b>Purification:</b> affinity purified by Protein A</p> <p><b>Concentration:</b> 1mg/ml</p> <p><b>Storage:</b> 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.</p> <p><b>Background:</b> This gene encodes a tyrosine kinase that binds Cdc42Hs in its GTP-bound form and inhibits both the intrinsic and GTPase-activating protein (GAP)-stimulated GTPase activity of Cdc42Hs. This binding is mediated by a unique sequence of 47 amino acids C-terminal to an SH3 domain. The protein may be involved in a regulatory mechanism that sustains the GTP-bound active form of Cdc42Hs and which is directly linked to a tyrosine phosphorylation signal transduction pathway. Several alternatively spliced transcript variants have been identified from this gene, but the full-length nature of only two transcript variants has been determined. [provided by RefSeq].</p>	<p><b>Isotype:</b> IgG</p> <p><b>SWISS:</b> Q07912</p> <p><b>Applications:</b> <b>WB</b> (1:500-2000) <b>IHC-P</b> (1:100-500) <b>IHC-F</b> (1:100-500) <b>IF</b> (1:100-500) <b>ELISA</b> (1:5000-10000)</p> <p><b>Reactivity:</b> (predicted: Human, Mouse, Rat, Pig, Cow, Dog)</p> <p><b>Predicted MW.:</b> 117 kDa</p> <p><b>Subcellular Location:</b> Cell membrane ,Cytoplasm ,Nucleus</p>
---	--