

bs-9665R**[Primary Antibody]****MREG Rabbit pAb**

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —

Host: Rabbit	Isotype: IgG	Applications: WB (1:500-2000) IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) ICC/IF (1:100-500) ELISA (1:5000-10000) Reactivity: (predicted: Human, Mouse, Rat) Predicted MW.: 25 kDa Subcellular Location: Cell membrane
Clonality: Polyclonal		
GeneID: 55686	SWISS: Q8N565	
Target: MREG		
Immunogen: KLH conjugated synthetic peptide derived from human DSU/MREG: 121-214/214.		
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: The photoreceptor rod cell that is responsible for vision under conditions of low light consists of stacked arrays of disk membranes that make up its outer segment portion. Regulated by complex biochemical mechanisms, the rod outer segment is under constant renewal as new disks form at the base. MREG is thought to play a role in membrane fusion and in regulating the biogenesis of disk membranes of photoreceptor rods. MREG interacts with RDS (also known as Peripherin-2), a photoreceptor specific tetraspanin protein that is required to maintain normal cell structure during the renewal process of membrane fusion. MREG is 214 amino acids in length, is expressed in photoreceptor cells and and is expressed as two isoforms due to alternative splicing.		