bsm-52235R

[Primary Antibody]

STAT3 Recombinant Rabbit mAb



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

DATASHEET -

Host: Rabbit Isotype: IgG Clonality: Recombinant CloneNo.: 6G5 **GeneID: 6774 SWISS:** P40763

Target: STAT3

Immunogen: A synthesized peptide derived from human STAT3: 1-33.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: The protein encoded by this gene is a member of the STAT protein family. In response to cytokines and growth factors, STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. This protein is activated through phosphorylation in response to various cytokines and growth factors including IFNs, EGF, IL5, IL6, HGF, LIF and BMP2. This protein mediates the expression of a variety of genes in response to cell stimuli, and thus plays a key role in many cellular processes such as cell growth and apoptosis. The small GTPase Rac1 has been shown to bind and regulate the activity of this protein. PIAS3 protein is a specific inhibitor of this protein. Mutations in this gene are associated with infantile-onset multisystem autoimmune disease and hyper-immunoglobulin E syndrome. Alternative splicing results in multiple transcript variants encoding distinct isoforms. [provided by RefSeq, Sep 2015]

Applications: WB (1:1000-5000)


IHC-P (1:100-500) **IHC-F** (1:100-500) **IF** (1:100-500) Flow-Cyt (1:50-100) ICC/IF (1:50-200)

Reactivity: Human, Mouse, Rat


Predicted MW.: 88 kDa

Subcellular Location: Cytoplasm ,Nucleus


- VALIDATION IMAGES -




25 ug total protein per lane of various lysates (see on figure) probed with STAT3 monoclonal antibody, unconjugated (bsm-52235R) at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at rt for 60 min




25 ug total protein per lane of various lysates (see on figure) probed with STAT3 monoclonal antibody, unconjugated (bsm-52235R) at 1:5000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at rt for 60 min



Paraformaldehyde-fixed, paraffin embedded Human Kidney; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with STAT3 Monoclonal Antibody, Unconjugated(bsm-52235R) at 1:200 overnight at 4°C, followed by conjugation to the bs-0295G-HRP and DAB (C-0010) staining.



Paraformaldehyde-fixed, paraffin embedded Human Kidney Cancer; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15



The Hela (H) cells were fixed with 4% PFA (10 min at r.t.) and then permeabilized with 90% icecold methanol for 20 min at -20°C, the cells then

min; Antibody incubation with STAT3
Monoclonal Antibody,
Unconjugated(bsm-52235R) at 1:200 overnight
at 4°C, followed by conjugation to the bs-0295GHRP and DAB (C-0010) staining.

were incubated in 5%BSA to block non-specific protein-protein interactions (30 min at r.t.). Primary Antibody (green): Rabbit Anti-STAT3 antibody (bsm-52235R): 1:50-100/10^6 cells; Secondary Antibody (white blue): Goat anti-Rabbit IgG-BF488 (bs-60295G-BF488): 1 µg/test. Isotype Control (orange): Rabbit IgG (bs-0295P). Blank control (black): PBS. Acquisition of 20,000 events was performed.

- SELECTED CITATIONS -

- [IF=8.2] Qingming Bao. et al. Tumor targeted siRNA delivery by adenosine receptor-specific curdlan nanoparticles. INT J BIOL MACROMOL. 2023 Dec;253:126845 WB ;Mouse. 37703972
- [IF=5.696] Jiang-Tao Fan. et al. Exosomal lncRNA NEAT1 from cancer-associated fibroblasts facilitates endometrial cancer progression via miR-26a/b-5p-mediated STAT3/YKL-40 signaling pathway. Neoplasia. 2021 Jul;23:692 WB;Human. 34153644
- [IF=5.195] Yong Wang. et al. An integrated network pharmacology approach reveals that Darutigenol reduces inflammation and cartilage degradation in a mouse collagen-induced arthritis model by inhibiting the JAK-STAT3 pathway. J ETHNOPHARMACOL. 2023 May;:116574 WB; Mouse. 37160212
- [IF=3.8] Junyu Chen. et al. Overexpression of FOS enhances the malignant potential of eutopic endometrial stromal cells in patients with endometriosis-associated ovarian cancer. oncol rep. 2025 Apr; 53(4):45. Western Blot; Human. 39981914
- [IF=2.57] Yuxi Jia. et al. Effects of Dietary n-3/n-6 PUFA Ratios on Animal Behaviors and Inflammation in a Prenatal Valproic Acid Exposure-Induced Rat Model of Autism. Neuropsych Dis Treat. 2022 Feb;18:401-412 WB;Rat. 10.2147/NDT.S343229