bs-25346R

[Primary Antibody]

Bioss ANTIBODIES

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

Mesothelin Rabbit pAb

- DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

GeneID: 10232 **SWISS:** Q13421

Target: Mesothelin

Immunogen: KLH conjugated synthetic peptide derived from human Mesothelin:

351-450/630.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: Mesothelin(MSLN)may play a role in cell adhesion and cell shape

dynamics; human homolog highly expressed in renal cell carcinoma. malignant tumors, such as a skin lump, ovary cancer and the pancreas cancer…etc..Mainly used for the research of the

malignant tumor.

Applications: IHC-P (1:100-500)


IHC-F (1:100-500) IF (1:100-500)

Reactivity: Human, Mouse, Rat


Predicted MW.: 64 kDa

Subcellular Secreted ,Cell membrane

VALIDATION IMAGES



Paraformaldehyde-fixed, paraffin embedded (human gastric carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Mesothelin) Polyclonal Antibody, Unconjugated (bs-25346R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructionsand DAB staining.



Paraformaldehyde-fixed, paraffin embedded (human tonsil); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Mesothelin) Polyclonal Antibody, Unconjugated (bs-25346R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructionsand DAB staining.