

bs-9522R**[Primary Antibody]****MPP1 Rabbit pAb****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

— DATASHEET —

| | |
|---|---|
| Host: Rabbit Clonality: Polyclonal GeneID: 4354 Target: MPP1 Immunogen: KLH conjugated synthetic peptide derived from human MPP1/EMP55: 101-200/466. Purification: affinity purified by Protein A Concentration: 1mg/ml Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. Background: The Kinesins constitute a large family of microtubule-dependent motor proteins which are responsible for the distribution of numerous organelles, vesicles and macromolecular complexes throughout the cell. Individual Kinesin members play crucial roles in cell division, intracellular transport and membrane trafficking events, including endocytosis and transcytosis. MPP1 (M-phase phosphoprotein 1), also known as KIF20B (kinesin family member 20B), MPHOSPH1 or KRMP1, is a 1,820 amino acid protein that localizes to both the nucleus and the cytoplasm and contains one kinesin-motor domain. Expressed in kidney, brain, testis and ovary, MPP1 functions as a plus-end directed motor enzyme that interacts with Pin1 and is required for the completion of cytokinesis. MPP1, which exists as multiple alternatively spliced isoforms termed 1-5, is subject to post-translational phosphorylation, probably by ATM or ATR. | Isotype: IgG SWISS: Q00013 Applications: WB (1:500-2000) Reactivity: Mouse (predicted: Human, Rat, Pig, Sheep, Cow, Dog) Predicted MW.: 52 kDa Subcellular Location: Cell membrane |
|---|---|

— VALIDATION IMAGES —

Sample: Liver (Mouse) Lysate at 40 ug Primary: Anti- MPP1 (bs-9522R) at 1/1000 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 52 kD
Observed band size: 52 kD



Sample: Spinal cord (Mouse) Lysate at 40 ug Primary: Anti- MPP1 (bs-9522R) at 1/1000 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 52 kD
Observed band size: 52 kD