

bs-16872R**[Primary Antibody]****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

ODCp Rabbit pAb**— DATASHEET —**


Host: Rabbit	Isotype: IgG	Applications: WB (1:500-2000) IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500)
Clonality: Polyclonal		
GeneID: 113451	SWISS: Q96A70	
Target: ODCp		
Immunogen: KLH conjugated synthetic peptide derived from human ODCp: 51-150/460.		
Purification: affinity purified by Protein A		Reactivity: Human, Mouse, Rat (predicted: Rabbit, Pig, Sheep, Cow, Dog, Horse)
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: ODCp is a 460 amino acid protein that is expressed in brain and testis. ODCp is a member of the Orn/Lys/Arg decarboxylase class-II family that catalyzes the creation of CO ₂ and agmatine from L-arginine. Mammalian ODCp differs from the forms expressed in bacteria and plants and shares less than 50% homology with Ornithine decarboxylase (ODC). ODCp is associated with the mitochondrial membrane where excess agmatine can be degraded by the enzyme Agmatinase or bound by the imidazoline recepto. In the brain, the highest levels of ODCp are found in the hypothalamus. Mammalian ODCp is thermally unstable and can be inhibited by Ca ²⁺ , Co ²⁺ and polyamines.		
		Predicted MW.: 50 kDa
		Subcellular Location: Cell membrane ,Cytoplasm ,Nucleus

— VALIDATION IMAGES —

Sample: Testis (Mouse) Lysate at 40 ug Primary:
Anti-ODCp(bs-16872R) at 1/300 dilution
Secondary: IRDye800CW Goat Anti-RabbitIgG at
1/20000 dilution Predicted band size: 50kD
Observed band size: 50kD



Sample: Raji (Human) Cell Lysate at 40 ug
Primary: Anti-ODCp(bs-16872R) at 1/300 dilution
Secondary: IRDye800CW Goat Anti-RabbitIgG at
1/20000 dilution Predicted band size: 50kD
Observed band size: 50kD



Paraformaldehyde-fixed, paraffin embedded (Rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (ODCp) Polyclonal Antibody, Unconjugated (bs-16872R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.