

bs-7602R**[Primary Antibody]****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

TIAF1 Rabbit pAb**— DATASHEET —**

Host: Rabbit	Isotype: IgG	Applications: IHC-P (1:100-500)
Clonality: Polyclonal		IHC-F (1:100-500)
GeneID: 9220	SWISS: Q95411	IF (1:100-500)
Target: TIAF1		Reactivity: Human
Immunogen: KLH conjugated synthetic peptide derived from human TIAF1: 34-115/115.		
Purification: affinity purified by Protein A		Predicted MW.: 12 kDa
Concentration: 1mg/ml		Subcellular Location: Nucleus
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: TIAF-1 (TGF Beta 1-induced antiapoptotic factor 1), also known as MYO18A, MAJN or MYSPDZ, is a TGF Beta 1-induced anti-apoptotic factor that localizes to the nucleus and functions to inhibit the cytotoxic effects of TNF? TRADD and FADD. Two isoforms of mouse TIAF-1 exist and are designated MysPDZ-Alpha and MysPDZ-Beta. MysPDZ-Alpha consists of a KE-rich region, an N-terminal PDZ domain and a prevalent myosin homologous head region, while the MysPDZ-Beta isoform lacks the KE-rich region and PDZ domain. MysPDZ-Alpha is present in most tissues and is known to co-localize with the ER-Golgi complex and with membrane ruffles and filopodia. MysPDZ-Beta is expressed specifically in hematopoietic tissues and cell lines and shows dispersed localization in the cytoplasm.		

— VALIDATION IMAGES —

Tissue/cell: human kidney tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Incubation: Anti-TIAF1 Polyclonal Antibody, Unconjugated(bs-7602R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining