

**bs-15523R****[ Primary Antibody ]****phospho-IFNGR1 (Tyr457) Rabbit pAb****Bioss**  
**ANTIBODIES**

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

**— DATASHEET —**


<b>Host:</b> Rabbit	<b>Isotype:</b> IgG	<b>Applications:</b> <b>WB</b> (1:500-2000) <b>IHC-P</b> (1:100-500) <b>IHC-F</b> (1:100-500) <b>IF</b> (1:100-500)
<b>Clonality:</b> Polyclonal		
<b>GeneID:</b> 3459	<b>SWISS:</b> P15260	
<b>Target:</b> phospho-IFNGR1 (Tyr457)		
<b>Immunogen:</b> KLH conjugated synthesised phosphopeptide derived from human IFNGR1 around the phosphorylation site of Tyr457: EF(p-S)-NH <sub>2</sub> .		
<b>Purification:</b> affinity purified by Protein A		<b>Reactivity:</b> Human, Mouse
<b>Concentration:</b> 1mg/ml		
<b>Storage:</b> 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		<b>Predicted MW.:</b> 53 kDa
<b>Background:</b> This gene (IFNGR1) encodes the ligand-binding chain(alpha) of the gamma interferon receptor. Human interferon-gamma receptor is a heterodimer of IFNGR1 and IFNGR2. A genetic variation in IFNGR1 is associated with susceptibility to Helicobacter pylori infection. In addition, defects in IFNGR1 are a cause of mendelian susceptibility to mycobacterial disease, also known as familial disseminated atypical mycobacterial infection. [provided by RefSeq].		<b>Subcellular Location:</b> Cell membrane

**— VALIDATION IMAGES —**

Sample: Jurkat (Human) Cell Lysate at 40 ug  
 Primary: Anti-p-IFNGR1 (bs-15523R) at 1/300 dilution  
 Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution  
 Predicted band size: 53 kD  
 Observed band size: 53 kD



Sample: 293T Cell (Human) Lysate at 40 ug  
 Primary: Anti-phospho-IFNGR1 (Tyr457) (bs-15523R) at 1/300 dilution  
 Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution  
 Predicted band size: 53 kD  
 Observed band size: 53 kD



Paraformaldehyde-fixed, paraffin embedded (mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Phospho-IFNGR1) Polyclonal Antibody, Unconjugated (bs-15523R) at 1:500 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.