


bsm-52225R**[Primary Antibody]****BioSS**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

Smad4 Recombinant Rabbit mAb**— DATASHEET —****Host:** Rabbit**Isotype:** IgG**Clonality:** Recombinant**CloneNo.:** 3A1**GeneID:** 4089**SWISS:** Q13485**Target:** Smad4**Immunogen:** A synthesized peptide derived from human SMAD4: 500-552/552.**Purification:** affinity purified by Protein A**Concentration:** 1mg/ml**Storage:** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.**Background:** Smad 4 is a member of the Mothers Against Dpp (MAD)-related family of proteins. So far, eight Smads have been identified and can be divided in 3 subgroups based on their structure and functions; pathway-restricted, common mediator and inhibitory Smad. Smad 4 is the common Smad (co-Smad). Previously identified as the tumor suppressor DPC4 (deleted in pancreatic carcinoma, locus 4), Smad 4 is functionally distinct among the Smad family, and is required for the assembly and transcriptional activation of diverse, Smad-DNA complexes. In contrast to the R-Smads, Smad 4 is not regulated by phosphorylation, but acts as a common mediator of TGF-Beta, activin, and bone morphogenetic protein signaling responses. Smad 4 is frequently inactivated in pancreatic, biliary and colorectal tumors.**Applications:** WB (1:500-2000)**IHC-P** (1:50-200)**IHC-F** (1:50-200)**IF** (1:50-200)**Reactivity:** Human, Mouse, Rat**Predicted
MW.:** 60 kDa**Subcellular
Location:** Cytoplasm ,Nucleus**— VALIDATION IMAGES —**

25 ug total protein per lane of various lysates (see on figure) probed with Smad4 monoclonal antibody, unconjugated (bsm-52225R) at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.



Paraformaldehyde-fixed, paraffin embedded Human Breast; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with Smad4 Monoclonal Antibody, Unconjugated(bsm-52225R) at 1:200 overnight at 4°C, followed by conjugation to the bs-0295G-HRP and DAB (C-0010) staining.



Paraformaldehyde-fixed, paraffin embedded Human Colon; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with Smad4 Monoclonal Antibody, Unconjugated(bsm-52225R) at 1:200 overnight at 4°C, followed by conjugation to the bs-0295G-HRP and DAB (C-0010) staining.