

bs-3067R**[Primary Antibody]****phospho-Bcr (Tyr177) Rabbit pAb**

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —

Host: Rabbit	Isotype: IgG	Applications: WB (1:500-2000) IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) ELISA (1:5000-10000) Reactivity: (predicted: Human, Mouse, Rat, Rabbit, Pig, GuineaPig) Predicted MW.: 143 kDa Subcellular Location: Cell membrane
Clonality: Polyclonal		
GeneID: 613	SWISS: P11274	
Target: Bcr (Tyr177)		
Immunogen: KLH conjugated Synthesisedphosphopeptide derived from human Bcr around the phosphorylation site of Tyr177: PF(p-Y)VN.		
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: A reciprocal translocation between chromosomes 22 and 9 produces the Philadelphia chromosome, which is often found in patients with chronic myelogenous leukemia. The chromosome 22 breakpoint for this translocation is located within the BCR gene. The translocation produces a fusion protein which is encoded by sequence from both BCR and ABL, the gene at the chromosome 9 breakpoint. Although the BCR-ABL fusion protein has been extensively studied, the function of the normal BCR gene product is not clear. The protein has serine/threonine kinase activity and is a GTPase-activating protein for p21rac. Two transcript variants encoding different isoforms have been found for this gene.		