

**bs-1941R****[ Primary Antibody ]****Zfp91 Rabbit pAb****BioSS**  
**ANTIBODIES**

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

**— DATASHEET —**

<b>Host:</b> Rabbit	<b>Isotype:</b> IgG	<b>Applications:</b> IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500)  <b>Reactivity:</b> Mouse, Rat (predicted: Human, Pig, Sheep, Cow, Chicken, Dog, Horse)  <b>Predicted MW.:</b> 63 kDa  <b>Subcellular Location:</b> Nucleus
<b>Clonality:</b> Polyclonal		
<b>GeneID:</b> 80829	<b>SWISS:</b> Q96JP5	
<b>Target:</b> Zfp91		
<b>Immunogen:</b> KLH conjugated synthetic peptide derived from human Zfp91: 301-400/570.		
<b>Purification:</b> affinity purified by Protein A		
<b>Concentration:</b> 1mg/ml		
<b>Storage:</b> 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
<b>Background:</b> The protein encoded by ZFP91 is a member of the zinc finger family of proteins. This protein contains C2H2 type domains, which are the classical zinc finger domains found in numerous nucleic acid- binding proteins. May be involved in transcriptional regulation. May play an important role in cell proliferation and/or anti- apoptosis Found in all the examined tissues including brain, heart, kidney, lung, liver, spleen, thymus, skeletal muscle, ovary and testis. Belongs to the krueppel C2H2-type zinc-finger protein family.		

**— VALIDATION IMAGES —**

Paraformaldehyde-fixed, paraffin embedded (Mouse testis); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Zfp91) Polyclonal Antibody, Unconjugated (bs-1941R) at 1:400 overnight at 4°C, followed by a conjugated secondary antibody (sp-0023) for 20 minutes and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Rat testis); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Zfp91) Polyclonal Antibody, Unconjugated (bs-1941R) at 1:400 overnight at 4°C, followed by a conjugated secondary antibody (sp-0023) for 20 minutes and DAB staining.