

bs-1912R**[Primary Antibody]****CGA Rabbit pAb****BioSS**
ANTIBODIES

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —

Host: Rabbit	Isotype: IgG	Applications: IHC-P (1:100-500)
Clonality: Polyclonal		IHC-F (1:100-500)
GeneID: 1081	SWISS: P01215	IF (1:100-500)
Target: CGA		ELISA (1:5000-10000)
Immunogen: Natural human hCG protein: 25-116/116.		
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		Reactivity: (predicted: Human)
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		Predicted MW.: 13 kDa
Background: The four human glycoprotein hormones chorionic gonadotropin (CG), luteinizing hormone (LH), follicle stimulating hormone (FSH), and thyroid stimulating hormone (TSH) are dimers consisting of alpha and beta subunits that are associated noncovalently. The alpha subunits of these hormones are identical, however, their beta chains are unique and confer biological specificity. The protein encoded by this gene is the alpha subunit and belongs to the glycoprotein hormones alpha chain family. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Nov 2011]		Subcellular Location: Secreted

— SELECTED CITATIONS —

- **[IF=12.7]** Dottermusch Matthias. et al. Pituitary neuroendocrine tumors with PIT1/SF1 co-expression show distinct clinicopathological and molecular features. ACTA NEUROPATHOL. 2024 Jun;147(1):1-13 IHC ;Human. 38228887