## bs-1926R

# [ Primary Antibody ]

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

# **UCP-2 Rabbit pAb**

DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Target: UCP-2

**Immunogen:** KLH conjugated synthetic peptide derived from mouse UCP-2:

201-309/309.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

**Background:** UCPs facilitate the transfer of anions from the inner to the outer

mitochondrial membrane and the return transfer of protons from the outer to the inner mitochondrial membrane. They also reduce the mitochondrial membrane potential in mammalian cells. UCP2 gene is expressed in many tissues, with the greatest expression in skeletal muscle. UCP2 is thought to play a role in non shivering

thermogenesis, obesity and diabetes.

Applications: WB (1:500-2000)

**IHC-P** (1:100-500) IHC-F (1:100-500) **IF** (1:100-500) Flow-Cyt (1µg /test)

ICC/IF (1:100-500)

Reactivity: Human, Mouse, Rat


(predicted: Rabbit, Pig,

Horse)


Predicted 34 kDa

Subcellular Location: Cytoplasm


## VALIDATION IMAGES




Sample: Lane 1: Stomach (Rat) Lysate at 40 ug Lane 2: NIH/3T3 (Mouse) Cell Lysate at 30 ug Lane 3: SiHa (Human) Cell Lysate at 30 ug Lane 4: DU145 (Human) Cell Lysate at 30 ug Lane 5: RAW264.7 (Mouse) Cell Lysate at 30 ug Lane 6: Liver (Mouse) Lysate at 40 ug Primary: Anti-UCP-2 (bs-1926R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 33 kD Observed band size: 33 kD




Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (UCP-2) Polyclonal Antibody, Unconjugated (bs-1926R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Hela cell; 4% Paraformaldehyde-fixed; Triton X-100 at room temperature for 20 min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min: Antibody incubation with (UCP-2) polyclonal Antibody, Unconjugated (bs-1926R) 1:100, 90 minutes at 37°C; followed by a conjugated Goat Anti-Rabbit IgG antibody at 37°C for 90 minutes, DAPI (blue, C02-04002) was used to stain the cell nuclei.



Hela cell; 4% Paraformaldehyde-fixed; Triton X-100 at room temperature for 20 min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Antibody incubation with (UCP-2) polyclonal Antibody, Unconjugated (bs-1926R) 1:100, 90 minutes at 37°C; followed by a conjugated Goat Anti-Rabbit IgG antibody at 37°C for 90 minutes, DAPI (blue, C02-04002) was



Blank control (blue line): Hela (blue). Primary Antibody (green line): Rabbit Anti-UCP-2 antibody (bs-1926R) Dilution: 1µg/10^6 cells; Isotype Control Antibody (orange line): Rabbit IgG . Secondary Antibody (white blue line): F(ab') 2 fragment goat anti-rabbit IgG-FITC Dilution: 1µg /test. Protocol The cells were fixed with 2% paraformaldehyde (10 min), then

used to stain the cell nuclei.

permeabilized with 90% ice-cold methanol for 30 min on ice.Cells stained with Primary Antibody for 30 min at room temperature. The cells were then incubated in 1 X PBS/2%BSA/10% goat serum to block nonspecific protein-protein interactions followed by the antibody for 15 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.

### - SELECTED CITATIONS -

- [IF=5.988] Lei Zhao. et al. Polysaccharides From Pogostemon cablin (Blanco) Benth.: Characterization and Antioxidant Activities. FRONT PHARMACOL. 2022; 13: 933669 WB; Mouse. 35784681
- [IF=5.469] Tidwell, Tia R.. et al. Metabolic flux analysis of 3D spheroids reveals significant differences in glucose metabolism from matched 2D cultures of colorectal cancer and pancreatic ductal adenocarcinoma cell lines. CANCER METAB. 2022 Dec;10(1):1-16 FCM; Human. 35578327
- [IF=4.35] Chuang, Yao-Chung, et al. "Peroxisome proliferator activated receptors γ/mitochondrial uncoupling protein 2 signaling protects against seizure-induced neuronal cell death in the hippocampus following experimental status Other ;Rat. 22849356
- [IF=4.35] Songsong Jiang. et al. A Comparison Study on the Therapeutic Effect of High Protein Diets Based on Pork Protein versus Soybean Protein on Obese Mice. FOODS. 2022 Jan;11(9):1227 WB; Mouse. 35563950
- [IF=2.6] Yaning Biao. et al. Wulingsan Alleviates MAFLD by Activating Autophagy via Regulating the AMPK/mTOR/ULK1 Signaling Pathway. CAN J GASTROENTEROL. 2024 Jul;2024(1):9777866 IHC; Rat. 39035827