bs-13856R

[Primary Antibody]

Bioss ANTIBODIES

www.bioss.com.cn sales@bioss.com.cl

sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

- DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

CEP192 Rabbit pAb

GenelD: 55125 **SWISS:** Q8TEP8

Target: CEP192

Immunogen: KLH conjugated synthetic peptide derived from human CEP192:

51-150/2537.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: Required for mitotic centrosome and spindle assembly. Appears to

be a major regulator of pericentriolar material (PCM) recruitment,

centrosome maturation, and centriole duplication.

Applications: IHC-P (1:100-500)


IHC-F (1:100-500) IF (1:100-500)

Reactivity: Human (predicted: Mouse)

Predicted 279 kDa

Subcellular Nucleus

VALIDATION IMAGES -



Tissue/cell: human liver cancer; 4%
Paraformaldehyde-fixed and paraffinembedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min; Incubation: Anti-CEP192 Polyclonal Antibody, Unconjugated(bs-13856R) 1:500, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining