

bs-0365R-HRP

[Secondary Antibodies]

Bioss
ANTIBODIES

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

human IgG Fc, HRP conjugated


— DATASHEET —

Host: Rabbit	Isotype: IgG	Applications: WB
Clonality: Polyclonal		IHC-P (1:100-500)
Target: human IgG Fc		IHC-F
Concentration: 1mg/ml		ICC/IF
Storage: 0.01M PBS(pH7.4). Shipped at 4°C. Store at -20 °C for one year. Avoid repeated freeze/thaw cycles.		ELISA
Background: The Fc region is the constant region on an immunoglobulin molecule, the area that is exactly the same on all antibodies. The Fc region is found on the heavy chains and is not involved in binding antigens.		Reactivity: Human
		Subcellular Location: Secreted


— VALIDATION IMAGES —



Sample: Lane 1: Human Serum Primary: Anti-human IgG Fc (bs-0365R) at 1/1000 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: kDa
Observed band size: 50 kDa



Paraformaldehyde-fixed, paraffin embedded (human colon); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (human IgG Fc) Polyclonal Antibody, Unconjugated (bs-0365R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (human colon carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (human IgG Fc) Polyclonal Antibody, Unconjugated (bs-0365R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.