bs-3123R

[Primary Antibody]

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

phospho-Cyclin B1 (Ser133) Rabbit pAb

DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

GeneID: 891 SWISS: P14635

Target: Cyclin B1 (Ser133)

Immunogen: KLH conjugated Synthesised phosphopeptide derived from human

Cyclin B1 around the phosphorylation site of Ser133: ET(p-S)GC.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: The protein encoded by this gene is a regulatory protein involved

in mitosis. The gene product complexes with p34(cdc2) to form the maturation-promoting factor (MPF). Two alternative transcripts have been found, a constitutively expressed transcript and a cell cycle-regulated transcript, that is expressed predominantly during G2/M phase. The different transcripts result from the use of alternate transcription initiation sites. [provided by RefSeq, Jul

2008].

Applications: Flow-Cyt (1µg /Test)

Reactivity: Human (predicted: Mouse,

ICC/IF (1:100-500)


Rat, Pig)

Predicted 48 kDa


MW.:

Subcellular Cytoplasm ,Nucleus

VALIDATION IMAGES -



Hela cell: 4% Paraformaldehyde-fixed: Triton X-100 at room temperature for 20 min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Antibody incubation with (Phospho-Cyclin B1 (Ser133)) polyclonal Antibody, Unconjugated (bs-3123R) 1:100, 90 minutes at 37°C; followed by a conjugated Goat Anti-Rabbit IgG antibody at 37°C for 90 minutes, DAPI (blue, C02-04002) was used to stain the cell nuclei.



Blank control (blue line): A549 (blue). Primary Antibody (green line): Rabbit Anti-Phospho-Cyclin B1 (Ser133) antibody (bs-3123R) Dilution: 1μg/10^6 cells; Isotype Control Antibody (orange line): Rabbit IgG . Secondary Antibody (white blue line): F(ab') 2 fragment goat antirabbit IgG-FITC Dilution: 1µg /test. Protocol The cells were fixed with 2% paraformaldehyde (10 min) and then permeabilized with 0.1% PBS-Tween for 20 min at room temperature. Cells stained with Primary Antibody for 30 min at room temperature. The cells were then incubated in 1 X PBS/2%BSA/10% goat serum to block non-specific protein-protein interactions followed by the antibody for 15 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.