# bs-2133R

- DATASHEET -

# [ Primary Antibody ]

# AT2R2 Rabbit pAb

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

#### Applications: WB (1:500-2000)

Reactivity: Mouse, Rat (predicted: Human, Pig, Sheep, Dog, Horse)

Predicted 41 kDa MW.:

Subcellular Location: Cell membrane

Clonality: Polyclonal GenelD: 186

SWISS: P50052

Isotype: IgG

Target: AT2R2

Host: Rabbit

Immunogen: KLH conjugated synthetic peptide derived from human angiotensin II receptor type 2: 151-250/363. < Extracellular >

Purification: affinity purified by Protein A

### Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: Angiotensin II (Ang II) is an important physiological effector of blood pressure and volume regulation through vasoconstriction, aldosterone release, sodium uptake and thirst stimulation. Although Ang II interacts with two types of cell surface receptors, AT1 and AT2, most of the major cardiovascular effects seem to be mediated through AT1. Molecular cloning of the AT1 protein has shown it to be a member of the G protein-associated seven transmembrane protein receptor family. Ang II treatment of cells results in activation of several signal transduction pathways as evidenced by tyrosine phosphorylation of several proteins and induction of others. PLC?is phosphorylated after 30 seconds of treatment with Angiotensin II, indicating this as an early signal transduction event. Ang II treatment also stimulates phosphorylation of Shc, FAK and MAP kinases, and induces MKP-1, indicating stimulation of growth factor pathways. Ang II stimulation through AT1 has been shown to activate the JAK/Stat pathway involving a direct interaction between JAK2 and AT1 as demonstrated by coimmunoprecipitation. The AT1 receptor has no cytoplasmic kinase domain, but is able to function as a substrate for Src kinases and has several putative phosphorylation sites.

## — VALIDATION IMAGES



Sample: Lane 1: Mouse Uterus tissue lysates Lane 2: Mouse Liver tissue lysates Lane 3: Rat Uterus tissue lysates Primary: Anti-AT2R2 (bs-2133R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 41 kDa Observed band size: 50 kDa

# – SELECTED CITATIONS —

• [IF=15.1] Xin Zhang. et al. Behind the Indolent Facade: Uncovering the Molecular Features and Malignancy Potential in Lung Minimally Invasive Adenocarcinoma by Single-Cell Transcriptomics. Advanced Science. 2023 Nov;:2303753 IF

;Human. 37991139