

bs-7574R**[Primary Antibody]****BioSS**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

RRAS Rabbit pAb**— DATASHEET —**

Host: Rabbit Clonality: Polyclonal GeneID: 6237 Target: RRAS Immunogen: KLH conjugated synthetic peptide derived from human RRAS: 131-218/218. Purification: affinity purified by Protein A Concentration: 1mg/ml Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. Background: H-Ras, K-Ras and N-Ras represent the prototype members of a family of small G proteins that are frequently activated to an oncogenic state in a wide variety of human tumors. Activation is due to point mutations at either position 12 or 61 within their coding sequence. Such mutations cause these proteins to be constitutively converted to their active, rather than the inactive, GDP-bound state. The related human R-Ras gene was initially cloned by low stringency hybridization methods. The R-Ras protein has been shown to interact with the Bcl-2 gene product involved in a signaling pathway that intervenes with apoptosis. Positions 38 and 87 (analogous to positions 12 and 61 in H-Ras) mutants of R-Ras have been shown to be capable of activating oncogenic function. Data has been obtained indicating that R-Ras may exert its biological effect by means of modulating the activity of the Raf-1 kinase on its direct downstream effectors.	Isotype: IgG SWISS: P10301 Applications: IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) Reactivity: Human (predicted: Mouse, Rat, Pig, Sheep, Cow, Dog) Predicted MW.: 23 kDa Subcellular Location: Cell membrane ,Cytoplasm
--	---

— VALIDATION IMAGES —

Paraformaldehyde-fixed, paraffin embedded (Human breast carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (RRAS) Polyclonal Antibody, Unconjugated (bs-7574R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.