

bs-7581R**[Primary Antibody]****ZHX2 Rabbit pAb****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

— DATASHEET —


Host: Rabbit	Isotype: IgG	Applications: IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) ICC/IF (1:50) Reactivity: Human, Rat (predicted: Mouse, Pig, Cow, Chicken, Dog, Horse) Predicted MW.: 92 kDa Subcellular Location: Nucleus
Clonality: Polyclonal		
GeneID: 22882	SWISS: Q9Y6X8	
Target: ZHX2		
Immunogen: KLH conjugated synthetic peptide derived from human ZHX2: 451-550/837.		
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: The members of the zinc fingers and homeoboxes gene family are nuclear homodimeric transcriptional repressors that interact with the A subunit of nuclear factor-Y (NF-YA) and contain two C2H2-type zinc fingers and five homeobox DNA-binding domains. This gene encodes member 2 of this gene family. In addition to forming homodimers, this protein heterodimerizes with member 1 of the zinc fingers and homeoboxes family. [provided by RefSeq, Jul 2008]. [FUNCTION] Acts as a transcriptional repressor. [SUBUNIT] Forms homodimers. Also forms heterodimers with ZHX1 and ZHX3. Heterodimerization with ZHX1 is not necessary for repressor activity. Interacts with NFYA. [SUBCELLULAR LOCATION] Nucleus. [TISSUE SPECIFICITY] Ubiquitously expressed. [SIMILARITY] Belongs to the ZHX family.		

— VALIDATION IMAGES —

Paraformaldehyde-fixed, paraffin embedded (human brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Incubation with (ZHX2) Polyclonal Antibody, Unconjugated (bs-7581R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Rat breast); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (ZHX2) Polyclonal Antibody, Unconjugated (bs-7581R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



MCF-7 cell; 4% Paraformaldehyde-fixed; Triton X-100 at room temperature for 20 min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Antibody incubation with (ZHX2) polyclonal Antibody, Unconjugated (bs-7581R) 1:50, 90 minutes at 37°C; followed by a conjugated Goat Anti-Rabbit IgG antibody at 37°C for 90 minutes, DAPI (blue, C02-04002) was used to stain the cell nuclei.