

bsm-34118M

[Primary Antibody]

DNA-RNA Hybrid Mouse mAb

Bioss
ANTIBODIES

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —

Host: Mouse

Isotype: IgG2a

Clonality: Monoclonal

CloneNo.: S9.6

Target: DNA-RNA Hybrid

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: PBS with 0.02% Proclin300 and 50% Glycerol, pH7.4.
Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Applications: IHC-P (1:100-500)

IHC-F (1:100-500)


IF (1:100-500)

ICC/IF (1:500-200)


ELISA (1:5000-10000)

Reactivity: Species independent


— VALIDATION IMAGES —



Paraformaldehyde-fixed, paraffin embedded Human Testicles; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with DNA-RNA Hybrid Monoclonal Antibody, Unconjugated (bsm-34118M) at 1:200 overnight at 4°C, followed by conjugation to the bs-40296G-HRP and DAB (C-0010) staining.



Paraformaldehyde-fixed, paraffin embedded Human Lung Cancer; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with DNA-RNA Hybrid Monoclonal Antibody, Unconjugated (bsm-34118M) at 1:200 overnight at 4°C, followed by conjugation to the bs-40296G-HRP and DAB (C-0010) staining.



Hela cell; 4% Paraformaldehyde-fixed; Triton X-100 at room temperature for 20 min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Antibody incubation with (DNA-RNA Hybrid) monoclonal Antibody, Unconjugated (bsm-34118M) 1:100, 90 minutes at 37°C; followed by a conjugated Goat Anti-Mouse IgG antibody at 37°C for 90 minutes, DAPI (blue, C02-04002) was used to stain the cell nuclei.