bs-2885R

[Primary Antibody]

Bioss ANTIBODIES

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn

GAB2 Rabbit pAb

- DATASHEET -

Host: Rabbit **Isotype:** IgG

Clonality: Polyclonal

GenelD: 9846 SWISS: Q9UQC2

Target: GAB2

Immunogen: KLH conjugated synthetic peptide derived from human Gab2:

601-676/676.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: This gene is a member of the GRB2-associated binding protein

(GAB) gene family. These proteins contain pleckstrin homology (PH) domain, and bind SHP2 tyrosine phosphatase and GRB2 adapter protein. They act as adapters for transmitting various signals in response to stimuli through cytokine and growth factor receptors, and T- and B-cell antigen receptors. The protein

encoded by this gene is the principal activator of

phosphatidylinositol-3 kinase in response to activation of the high affinity IgE receptor. Two alternatively spliced transcripts encoding different isoforms have been described for this gene. [provided by

RefSeq, Nov 2009]

Applications: IHC-P (1:100-500)

400-901-9800

IHC-F (1:100-500) IF (1:100-500) Flow-Cyt (1µg /test)

Reactivity: Human, Mouse, Rat


(predicted: Rabbit, Pig, Cow, Dog, Horse)

Predicted MW.: ^{74 kDa}

Subcellular

Location: Cell membrane ,Cytoplasm


VALIDATION IMAGES



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by microwave in sodium citrate buffer (pH6.0); Block endogenous peroxidase by 3% hydrogen peroxide for 30 minutes; Blocking buffer (3% BSA) at RT for 30min; Antibody incubation with (GAB2) Polyclonal Antibody, Unconjugated (bs-2885R) at 1:400 overnight at 4°C, followed by conjugation to the secondary antibody (labeled with HRP) and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Rat brain); Antigen retrieval by microwave in sodium citrate buffer (pH6.0); Block endogenous peroxidase by 3% hydrogen peroxide for 30 minutes; Blocking buffer (3% BSA) at RT for 30min; Antibody incubation with (GAB2) Polyclonal Antibody, Unconjugated (bs-2885R) at 1:400 overnight at 4°C, followed by conjugation to the secondary antibody (labeled with HRP) and DAB staining.



Blank control(blue): U-87MG(fixed with 2% paraformaldehyde (10 min) , then permeabilized with 90% ice-cold methanol for 30 min on ice). Primary Antibody:Rabbit Anti-DAB2 antibody(bs-2885R), Dilution: $1\mu g$ in $100~\mu L$ 1X PBS containing 0.5% BSA; Isotype Control Antibody: Rabbit IgG(orange) ,used under the same conditions); Secondary Antibody: Goat anti-rabbit IgG-PE(white blue), Dilution: 1:200 in 1 X PBS containing 0.5% BSA.

- SELECTED CITATIONS -

- [IF=3.118] Fu S et al. Berberine suppresses mast cell-mediated allergic responses via regulating FcɛRI-mediated and MAPK signaling.Int Immunopharmacol. 2019 Jun;71:1-6. WB; Human. 30861392
- [IF=2.31] Zhan, Ao, et al. "Decreased expression of Gab2 in patients with temporal lobe epilepsy and pilocarpine induced rat model." Synapse (2013). IHC; Human. 24327320