

bs-2052R**[Primary Antibody]****Transferrin Rabbit pAb****BioSS**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

— DATASHEET —

Host: Rabbit Clonality: Polyclonal GeneID: 7018 Target: Transferrin Purification: affinity purified by Protein A Concentration: 1mg/ml Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. Background: This gene encodes a glycoprotein with an approximate molecular weight of 76.5 kDa. It is thought to have been created as a result of an ancient gene duplication event that led to generation of homologous C and N-terminal domains each of which binds one ion of ferric iron. The function of this protein is to transport iron from the intestine, reticuloendothelial system, and liver parenchymal cells to all proliferating cells in the body. This protein may also have a physiologic role as granulocyte/pollen-binding protein (GPBP) involved in the removal of certain organic matter and allergens from serum. [provided by RefSeq, Sep 2009].	Isotype: IgG SWISS: P02787	Applications: WB (1:2000-10000) Reactivity: Human, Mouse, Rat, Goat Predicted MW.: 77 kDa Subcellular Location: Secreted
---	---	---

— VALIDATION IMAGES —

25 ug total protein per lane of various lysates (see on figure) probed with Transferrin polyclonal antibody, unconjugated (bs-2052R) at 1:10000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.



Sample: Lane 1: Serum (Human) at 20 ug Lane 2: Serum (Mouse) at 20 ug Lane 3: Serum (Rat) at 20 ug Lane 4: Serum (Goat) at 20 ug Primary: Anti-Transferrin (bs-2052R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 77 kD Observed band size: 77 kD