


bs-1741R**[Primary Antibody]****alpha Actinin 4 Rabbit pAb****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

DATASHEET**Host:** Rabbit**Isotype:** IgG**Clonality:** Polyclonal**GeneID:** 81**SWISS:** O43707**Target:** alpha Actinin 4**Immunogen:** KLH conjugated synthetic peptide derived from human Actinin alpha 4: 811-911/911.**Purification:** affinity purified by Protein A**Concentration:** 1mg/ml**Storage:** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.**Background:** Alpha actinins belong to the spectrin gene superfamily which represents a diverse group of cytoskeletal proteins, including the alpha and beta spectrins and dystrophins. Alpha actinin is an actin-binding protein with multiple roles in different cell types. In nonmuscle cells, the cytoskeletal isoform is found along microfilament bundles and adherens-type junctions, where it is involved in binding actin to the membrane. In contrast, skeletal, cardiac, and smooth muscle isoforms are localized to the Z-disc and analogous dense bodies, where they help anchor the myofibrillar actin filaments. This gene encodes a nonmuscle, alpha actinin isoform which is concentrated in the cytoplasm, and thought to be involved in metastatic processes. Mutations in this gene have been associated with focal and segmental glomerulosclerosis. [provided by RefSeq, Jul 2008].**Applications:** WB (1:500-2000)**IHC-P** (1:100-500)**IHC-F** (1:100-500)**IF** (1:100-500)**Flow-Cyt** (2ug/Test)**Reactivity:** Human, Mouse, Rat
(predicted: Rabbit, Pig, Cow, Chicken)**Predicted MW.:** 105 kDa**Subcellular Location:** Cytoplasm ,Nucleus**VALIDATION IMAGES**


Sample: Lane 1: Mouse Uterus tissue lysates
Lane 2: Rat Uterus tissue lysates Primary: Anti-alpha Actinin 4 (bs-1741R) at 1/1000 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 105 kDa
Observed band size: 105 kDa



Paraformaldehyde-fixed, paraffin embedded (Human breast carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (alpha Actinin 4-Loading Control) Polyclonal Antibody, Unconjugated (bs-1741R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Tissue/cell: rat brain tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Incubation: Anti-alpha-Actinin-4 Polyclonal Antibody, Unconjugated(bs-1741R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Blank control (black line) :U251. Primary
Antibody (green line): Rabbit Anti-alpha Actinin 4
antibody (bs-1741R) Dilution:2ug/Test;
Secondary Antibody (white blue line) : Goat
anti-rabbit IgG-AF488 Dilution: 0.5ug/Test.
Isotype control (orange line) : Normal Rabbit
IgG Protocol The cells were fixed with 4% PFA
(10min at room temperature)and then
permeabilized with 90% ice-cold methanol for
20 min at -20°C, The cells were then incubated in
5%BSA to block non-specific protein-protein
interactions for 30 min at room temperature
.Cells stained with Primary Antibody for 30 min
at room temperature. The secondary antibody
used for 40 min at room temperature.
Acquisition of 20,000 events was performed.

— SELECTED CITATIONS —

- **[IF=9.965]** Qinghua Yu. et al. Dual-Emission ZAlSe/ZnS Quantum Dots for Multi-level Bio-Imaging: Foam Cells and Atherosclerotic Plaque Imaging. J COLLOID INTERF SCI. 2022 Aug;: WB ;Mouse. 36088689
- **[IF=3.5]** Jinkui Sun. et al. Effect of Bovine MEF2A Gene Expression on Proliferation and Apoptosis of Myoblast Cells. GENES-BASEL. 2023 Jul;14(7):1498 ICC ;Bovine. 37510401