

**bsm-60595R**

**[ Primary Antibody ]**

**Bioss**  
ANTIBODIES

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

## TFEB Recombinant Rabbit mAb

### DATASHEET

**Host:** Rabbit

**Clonality:** Recombinant

**Target:** TFEB

**Purification:** affinity purified by Protein A

**Concentration:** 1mg/ml

**Storage:** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.  
Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

**Background:** Transcription factors of the basic helix-loop-helix zipper (bHLH-ZIP) family contain a basic domain, used for DNA binding, and HLH and ZIP domains, both used for oligomerization. TFEB was isolated from a human B-cell cDNA library using a binding sequence from the adenovirus major late promoter. This transcription factor probably binds to the USF/MLTF site and probably recognizes E-box sequences in the heavy-chain immunoglobulin enhancer.

**Isotype:** IgG

**CloneNo.:** D11B4

**Applications:** WB (1:500-1000)

**IHC-P** (1:100-500)

**IHC-F** (1:100-500)

**IF** (1:100-500)


**Flow-Cyt** (1:50-100)

**ICC/IF** (1:50)


**Reactivity:** Human

**Subcellular Location:** Cytoplasm ,Nucleus


### VALIDATION IMAGES




Blocking buffer: 5% NFDN/TBST Primary ab dilution: 1:1000 Primary ab incubation condition: 2 hours at room temperature Secondary ab: Goat Anti-Rabbit IgG H&L (HRP) Lysate: 1: Raji, 2: HeLa, 3: HepG2, 4: K562 Protein loading quantity: 20 µg Exposure time: 60 s Predicted MW: 53 kDa Observed MW: 66 kDa




Tissue: Human prostatic hyperplasia Section type: Formalin fixed & Paraffin -embedded section Retrieval method: High temperature and high pressure Retrieval buffer: Tris/EDTA buffer, pH 9.0 Primary ab dilution: 1:1000 Primary ab incubation condition: 1 hour at room temperature Secondary ab: SP Kit(Rabbit) (sp-0023) Counter stain: Hematoxylin (Blue) Comment: Color brown is the positive signal for bsm-60595R



Tissue: Human breast cancer Section type: Formalin fixed & Paraffin -embedded section Retrieval method: High temperature and high pressure Retrieval buffer: Tris/EDTA buffer, pH 9.0 Primary ab dilution: 1:1000 Primary ab incubation condition: 1 hour at room temperature Secondary ab: SP Kit(Rabbit) (sp-0023) Counter stain: Hematoxylin (Blue) Comment: Color brown is the positive signal for bsm-60595R



Cell line: Raji Fixative: 4% Paraformaldehyde Permeabilization: 0.1% TritonX-100 Primary ab dilution: 1:50 Primary incubation condition: 4°C overnight Secondary ab: Goat Anti-Rabbit IgG Nuclear counter stain: DAPI (Blue) Comment: Color green is the positive signal for bsm-60595R



Cell line: Raji Fixative: 4% Paraformaldehyde Permeabilization: 90% methanol Primary ab dilution: 1:100 Secondary ab: Goat anti Rabbit IgG Unlabelled control: The cell without incubation with primary antibody and secondary antibody (Black line). Isotype control: Rabbit monoclonal IgG (Blue line). Comment: Line red is the positive signal for bsm-60595R