

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

CDK8 Recombinant Rabbit mAb

Catalog Number: bsm-60609R

Target Protein: CDK8
Concentration: 1mg/ml

Form: Liquid Host: Rabbit

Clonality: Recombinant

Clone No.: B7D5

Isotype: IgG

Applications: WB (1:500-1:1000), IHC-P (1:100-500), IHC-F (1:100-500), IF

Reactivity: Human, Mouse

Subcellular Nucleus

Locations:

Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: This gene encodes a member of the cyclin-dependent protein kinase (CDK) family. CDK


family members are known to be important regulators of cell cycle progression. This kinase

and its regulatory subunit, cyclin C, are components of the Mediator transcriptional regulatory complex, involved in both transcriptional activation and repression by


phosphorylation of the carboxy-terminal domain of the largest subunit of RNA polymerase II. This kinase regulates transcription by targeting the cyclin-dependent kinase 7 subunits of the general transcription initiation factor IIH, thus providing a link between the Mediator complex and the basal transcription machinery. Multiple pseudogenes of this gene have been identified. Alternative splicing results in multiple transcript variants. [provided by

RefSeq, Oct 2016][provided by RefSeq, Jul 2008].

VALIDATION IMAGES



Tissue: Mouse colon Section type: Formalin fixed & Paraffin -embedded section Retrieval method: High temperature and high pressure Retrieval buffer: Tris/EDTA buffer, pH 9.0 Primary ab dilution: 1:100 Primary ab incubation condition: 1 hour at room temperature Secondary ab: SP Kit(Rabbit) (sp-0023) Counter stain: Hematoxylin (Blue) Comment: Color brown is the positive signal for bsm-60609R



Tissue: Human breast cancer Section type: Formalin fixed & Paraffin -embedded section Retrieval method: High temperature and high pressure Retrieval buffer: Tris/EDTA buffer, pH 9.0 Primary ab dilution: 1:100 Primary ab incubation condition: 1 hour at room temperature Secondary ab: SP Kit(Rabbit) (sp-0023) Counter stain: Hematoxylin (Blue) Comment: Color brown is the positive signal for bsm-60609R