

bs-9568R**[Primary Antibody]****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

TNIP1 Rabbit pAb**— DATASHEET —**

Host: Rabbit	Isotype: IgG	Applications: IHC-P (1:100-500)
Clonality: Polyclonal		IHC-F (1:100-500)
GeneID: 10318	SWISS: Q15025	IF (1:100-500)
Target: TNIP1		Reactivity: Rat (predicted: Human, Mouse, Rabbit, Pig, Cow, Dog, Horse)
Immunogen: KLH conjugated synthetic peptide derived from human TNIP1: 351-450/636.		Predicted MW.: 72 kDa
Purification: affinity purified by Protein A		Subcellular Location: Cytoplasm ,Nucleus
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: Abin-1 is a 636 amino acid protein that localizes to both the nucleus and the cytoplasm and is shuttled between the two intercellular regions in a CRM1-dependent manner. Expressed ubiquitously with highest expression in spleen and skeletal muscle, Abin-1 interacts with A20 and, via this interaction, interferes with TRAF2-mediated transactivation signals and effectively inhibits TNF-induced NF- κ B expression. Additionally, Abin-1 can be incorporated into HIV-1 virions and, if overexpressed, can inhibit viral replication. Abin-1 may also play an important role in the regulation of nuclear import and export activities. Multiple isoforms of Abin-1 exist due to alternative splicing events.		

— VALIDATION IMAGES —

Paraformaldehyde-fixed, paraffin embedded (rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (TNIP1) Polyclonal Antibody, Unconjugated (bs-9568R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Tissue/cell: rat brain tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Incubation: Anti-TNIP1 Polyclonal Antibody, Unconjugated(bs-9568R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining

— SELECTED CITATIONS —

- **[IF=1.813]** Zhou Xueling, et al. A20-Binding Inhibitor of NF- κ B 1 Ameliorates Neuroinflammation and Mediates Antineuroinflammatory Effect of Electroacupuncture in Cerebral Ischemia/Reperfusion Rats. Evid-Based Compl Alt. 2020;2020:6980398 IF ;Rat. 33110436