bs-2104R

[Primary Antibody]



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

TNFAIP1 Rabbit pAb

- DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

GenelD: 7126 **SWISS:** Q13829

Target: TNFAIP1

Immunogen: KLH conjugated synthetic peptide derived from human TNFAIP1:

231-316/316.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: This gene was identified as a gene whose expression can be

induced by the tumor necrosis factor alpha (TNF) in umbilical vein endothelial cells. Studies of a similar gene in mouse suggest that the expression of this gene is developmentally regulated in a

tissue-specific manner. [provided by RefSeq, Jul 2008].

Applications: IHC-P (1:100-500)


IHC-F (1:100-500) **IF** (1:100-500)

Reactivity: Human, Mouse, Rat

Predicted MW.: 35 kDa

Subcellular Cytoplasm ,Nucleus

VALIDATION IMAGES



Tissue/cell: Rat pancreas tissue; 4%
Paraformaldehyde-fixed and paraffinembedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min; Incubation: Anti-TNFAIP1 Polyclonal Antibody, Unconjugated(bs-2104R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Paraformaldehyde-fixed, paraffin embedded (human rectal carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (TNFAIP1) Polyclonal Antibody, Unconjugated (bs-2104R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.