bs-3242R

[Primary Antibody]

phospho-c-Kit (Tyr721) Rabbit pAb



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

GenelD: 3815 **SWISS:** P10721

Target: c-Kit (Tyr721)

Immunogen: KLH conjugated Synthesised phosphopeptide derived from human

c-Kit around the phosphorylation site of Tyr721: NE(p-Y)MD.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: c-Kit is a transmembrane tyrosine kinase encoded by the cKit proto

oncogene. c-Kit acts to regulate a variety of biological responses including cell proliferation, apoptosis, chemotaxis and adhesion.

Ligand binding to the extracellular domain leads to autophosphorylation on several tyrosine residues within the cytoplasmic domain, and activation. Mutations in c-Kit have been found to be important for tumor growth and progression in a variety of cancers including mast cell diseases, gastrointestinal stromal tumor, acute myeloid leukemia, Ewing sarcoma and lung cancer. Phosphorylation at tyrosine 721 of c-Kit allows binding and

activation of PI3 kinase.

Applications: IHC-P (1:100-500)

IHC-F (1:100-500) **IF** (1:100-500) Flow-Cyt (1µg/Test)

Reactivity: Human, Mouse


(predicted: Rat)

Predicted


105 kDa MW.:

Subcellular Cell membrane ,Cytoplasm

VALIDATION IMAGES



Paraformaldehyde-fixed, paraffin embedded (mouse cerebellum tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min: Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (c-Kit(Tyr721)) Polyclonal Antibody, Unconjugated (bs-3242R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Molt-4 cells were fixed with 4% PFA for 10min at room temperature, permeabilized with 20% PBST for 20 min at room temperature, and incubated in 5% BSA blocking buffer for 30 min at room temperature. Cells were then stained with CD117 Antibody(bs-3242R)at 1:100 dilution in blocking buffer and incubated for 30 min at room temperature, washed twice with 2%BSA in PBS, followed by secondary antibody incubation for 40 min at room temperature. Acquisitions of 20,000 events were performed. Cells stained with primary antibody (green), and isotype control (orange).

- SELECTED CITATIONS -

- [IF=2.12] Halsey, C. H. C., et al. "Expression of Phosphorylated KIT in Canine Mast Cell Tumor." Veterinary Pathology (2017): 0300985816688943. WB; Dog. 28129097
- [IF=2.379] Thamm DH et al. Phosphorylated KIT as a Predictor of Outcome in Canine Mast Cell Tumours Treated with Toceranib Phosphate or Vinblastine. Vet Comp Oncol. 2019 Jul 31. IHC; Human. 31365175