

bs-3007R**[Primary Antibody]****phospho-ASK1 (Ser966) Rabbit pAb****BioSS**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

— DATASHEET —

Host: Rabbit Clonality: Polyclonal GeneID: 4217 Target: ASK1 (Ser966) Immunogen: KLH conjugated Synthesised phosphopeptide derived from human ASK1 around the phosphorylation site of Ser966: SI(p-S)LP. Purification: affinity purified by Protein A Concentration: 1mg/ml Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. Background: Mitogen-activated protein kinase (MAPK) signaling cascades include MAPK or extracellular signal-regulated kinase (ERK), MAPK kinase (MKK or MEK), and MAPK kinase kinase (MAPKKK or MEKK). MAPKK kinase/MEKK phosphorylates and activates its downstream protein kinase, MAPK kinase/MEK, which in turn activates MAPK. The kinases of these signaling cascades are highly conserved, and homologs exist in yeast, Drosophila, and mammalian cells. MAPKKK5 contains 1,374 amino acids with all 11 kinase subdomains. Northern blot analysis shows that MAPKKK5 transcript is abundantly expressed in human heart and pancreas. The MAPKKK5 protein phosphorylates and activates MKK4 (aliases SERK1, MAPKK4) in vitro, and activates c-Jun N-terminal kinase (JNK)/stress-activated protein kinase (SAPK) during transient expression in COS and 293 cells; MAPKKK5 does not activate MAPK/ERK.	Isotype: IgG SWISS: Q99683 Applications: IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) Reactivity: Human, Mouse, Rat (predicted: Chicken, Dog, Horse) Predicted MW.: 152 kDa Subcellular Location: Cytoplasm
---	--

— VALIDATION IMAGES —

Paraformaldehyde-fixed, paraffin embedded Human Colon Cancer; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with phospho-ASK1 (Ser966) Polyclonal Antibody, Unconjugated (bs-3007R) at 1:200 overnight at 4°C, followed by conjugation to the bs-0295G-HRP and DAB (C-0010) staining.



Paraformaldehyde-fixed, paraffin embedded (rat heart); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (phospho-ASK1 (Ser966)) Polyclonal Antibody, Unconjugated (bs-3007R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

— SELECTED CITATIONS —

- **[IF=6.312]** Sou Hyun Kim. et al. Downregulation of Glutathione-Mediated Detoxification Capacity by Binge Drinking Aggravates Acetaminophen-Induced Liver Injury through IRE1α and ER Stress Signaling. Antioxidants-Basel. 2021 Dec;10(12):1949 WB ;Mouse. 34943052

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

- **[IF=4.109]** Wenying Sun. et al. Eucalyptol antagonized the apoptosis and immune dysfunction of grass carp hepatocytes induced by tetrabromobisphenol A by regulating ROS/ASK1/JNK pathway. ENVIRON TOXICOL. 2023 Jan;; WB ;Fish. 36629057
- **[IF=4.4]** Zheng Xiao. et al. GS-4997 halts the progression of tubulointerstitial injury in lupus nephritis. FASEB J. 2024 Dec;38(24):e70253 IHC ;Mouse. 39680018
- **[IF=2.3]** Erika Seki. et al. ASK1 activation in glial cells in post-mortem multiple sclerosis tissue. NEUROPATHOLOGY. 2024 May;; IHC,IF ;Human. 38775061