



Recombinant human EPOR protein, C-His (HEK293)

Catalog Number: bs-43588P

Concentration: >0.5mg/ml

Species: Human

AA Seq: 25-250/508

Predicted MW: 25.9

Tags: C-His

Endotoxin: Not tested

Purity: >90% as determined by SDS-PAGE

Purification: AC

Storage: Stored at -70°C or -20°C. Avoid repeated freeze/thaw cycles.

Background: The erythropoietin receptor (EPOR) is a member of the cytokine receptor family. There are

several isoforms including: EPOR-F (full length), EPOR-S (soluble form), and EPOR-T (truncated form). Upon erythropoietin (EPO) binding, the EPOR activates Jak2 tyrosine kinase which activates different intracellular pathways including: Ras/MAP kinase,

phosphatidylinositol 3-kinase and STAT transcription factors. The stimulated EPOR appears to have a role in erythroid cell survival. Defects in the EPOR may produce erythroleukemia and familial erythrocytosis. A functional EPOR is found in the cardiovascular system, including endothelial cells and cardiomyocytes, and data suggest that the EPO/EPO receptor system plays an important role in cardiac function. In animal studies, treatment with EPO during ischemia/reperfusion in the heart has been shown to limit the infarct size

and the extent of apoptosis.

VALIDATION IMAGES



The purity of the protein is greater than 90% as determined by reducing SDS-PAGE.