

---

## CDC2L5 Rabbit pAb

Catalog Number: bs-7897R

Target Protein: CDC2L5

Concentration: 1mg/ml

Form: Liquid

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: IHC-P (1:100-500), IHC-F (1:100-500), IF (1:100-500), ELISA (1:5000-10000)

Reactivity: Rat (predicted:Human, Mouse, Pig, Sheep, Cow, Dog)

Predicted MW: 165 kDa

Subcellular Nucleus

Locations:

Entrez Gene: 8621

Source: KLH conjugated synthetic peptide derived from human CDC2L5: 601-700/1512.

Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

**Background:** Cyclin-dependent kinase which displays CTD kinase activity and is required for RNA splicing. Has CTD kinase activity by hyperphosphorylating the C-terminal heptapeptide repeat domain (CTD) of the largest RNA polymerase II subunit RPB1, thereby acting as a key regulator of transcription elongation. Required for RNA splicing, probably by phosphorylating SRSF1/SF2. Required during hematopoiesis. In case of infection by HIV-1 virus, interacts with HIV-1 Tat protein acetylated at 'Lys-50' and 'Lys-51', thereby increasing HIV-1 mRNA splicing and promoting the production of the doubly spliced HIV-1 protein Nef. Tissue specificity: Expressed in fetal brain, liver, muscle and in adult brain. Also expressed in neuroblastoma and glioblastoma tumors.