

**bsm-54202R****[ Primary Antibody ]****BioSS**  
**ANTIBODIES**

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

**Nuclear Matrix Protein p84 Recombinant Rabbit mAb****— DATASHEET —**

<b>Host:</b> Rabbit	<b>Isotype:</b> IgG	<b>Applications:</b> WB (1:500-1000)  <b>Reactivity:</b> Human, Mouse, Rat  <b>Predicted MW.:</b> 72 kDa  <b>Subcellular Location:</b> Cytoplasm ,Nucleus
<b>Clonality:</b> Recombinant	<b>CloneNo.:</b> 6C1	
<b>GeneID:</b> 9984	<b>SWISS:</b> Q96FV9	
<b>Target:</b> Nuclear Matrix Protein p84		
<b>Purification:</b> affinity purified by Protein A		
<b>Concentration:</b> 1mg/ml		
<b>Storage:</b> 1*TBS (pH7.4), 0.05% BSA, 40% Glycerol. Preservative: 0.02% Proclin300. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
<b>Background:</b> This protein is part of the THO complex (THOC). The THO complex is recruited to transcribed genes and travels with the RNA polymerase during elongation. It may physically link proteins that function in transcription and in RNA export and RNA processing.		

**— VALIDATION IMAGES —**

Sample: Lane 1: Mouse Testis tissue lysates Lane 2: Rat Testis tissue lysates Lane 3: Human U251 cell lysates Lane 4: Human HUVEC cell lysates Lane 5: Human LOVO cell lysates Primary: Anti-Nuclear Matrix Protein p84 (bsm-54202R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 72 kDa Observed band size: 76 kDa



Sample: Lane 1: Mouse Placenta tissue lysates Lane 2: Rat Placenta tissue lysates Lane 3: Human HeLa cell lysates Lane 4: Human HepG2 cell lysates Primary: Anti-Nuclear Matrix Protein p84 (bsm-54202R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 72 kDa Observed band size: 75 kDa