### bs-25683R

## [ Primary Antibody ]

# Bioss

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

# eNOS Rabbit pAb

- DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**GenelD:** 4846 **SWISS:** P29474

Target: eNOS

**Immunogen:** KLH conjugated synthetic peptide derived from human eNOS:

21-120/1203.

**Purification:** affinity purified by Protein A

Concentration: 1mg/ml

**Storage:** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

**Background:** Nitric oxide synthase NOS oxidizes a guanidine nitrogen of arginine

releasing nitric oxide in the form of a free radical and citrulline. Nitric oxide thus generated acts as a messenger in diverse functions including vasodilation neurotransmission, anti tumor and anti pathogenic activities. NOS is classified under three types: neuronal NOS (nNOS) or brain NOS (bNOS); inducible NOS (iNOS)

or macrophage NOS (mNOS); and endothelial NOS (eNOS).

Applications: IHC-P (1:100-500)

**IHC-F** (1:100-500) **IF** (1:100-500)

Reactivity: Human (predicted: Mouse,


Rat, Pig, Sheep, Cow, Dog)

Predicted MW.: 133 kDa


.....

**Subcellular Location:** Cell membrane ,Cytoplasm

#### VALIDATION IMAGES



Paraformaldehyde-fixed, paraffin embedded (Human kidney); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (eNOS) Polyclonal Antibody, Unconjugated (bs-25683R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (human lung carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (eNOS) Polyclonal Antibody, Unconjugated (bs-25683R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.