## bs-2468R

## [ Primary Antibody ]

## MCP4 Rabbit pAb



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

- DATASHEET		400-901-9800
Host: Rabbit	Isotype: IgG	Applications: IHC-P (1:100-500) IHC-F (1:100-500)
Clonality: Polyclonal		IF (1:100-500)
<b>GenelD:</b> 6357	<b>SWISS:</b> Q99616	<b>ELISA</b> (1:5000-10000)
Target: MCP4		Reactivity: (predicted: Human, Rabbit,
Immunogen: KLH conjugated synthetic peptide derived from human CCL13: 51-98/98.		Dog)
Purification: affinity purified by	Protein A	
Concentration: 1mg/ml		Predicted MW.: <sup>110</sup> kDa
<b>Storage:</b> 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		Subcellular Secreted
<b>Background:</b> This gene is one of several Cys-Cys (CC) cytokine genes clustered on the q-arm of chromosome 17. Cytokines are a family of secreted proteins involved in immunoregulatory and inflammatory processes. The CC cytokines are proteins characterized by two adjacent cysteines. The cytokine encoded by this gene displays chemotactic activity for monocytes, lymphocytes, basophils and eosinophils, but not neutrophils. This chemokine plays a role in accumulation of leukocytes during inflammation. It may also be involved in the recruitment of monocytes into the arterial wall during artherosclerosis. [provided by RefSeq].		d