bs-2545R

[Primary Antibody]

Bioss ANTIBODIES

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

IL-2R gamma Rabbit pAb

- DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

GenelD: 3561 **SWISS:** P31785

Target: IL-2R gamma

Immunogen: KLH conjugated synthetic peptide derived from human IL-2R

gamma: 51-150/369. < Extracellular >

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: The protein encoded by this gene is an important signaling component of many interleukin receptors, including those of

interleukin -2, -4, -7 and -21, and is thus referred to as the common gamma chain. Mutations in this gene cause X-linked severe combined immunodeficiency (XSCID), as well as X-linked combined immunodeficiency (XCID), a less severe

immunodeficiency disorder. [provided by RefSeq, Mar 2010]

Applications: Flow-Cyt $(1\mu g / test)$ ICC/IF (1:100-500)


·

Reactivity: Human (predicted: Mouse, Rat, Pig, Cow, Dog)


Predicted MW.: 41 kDa

Subcellular Location: Cell membrane


VALIDATION IMAGES




SHSY5Y cell; 4% Paraformaldehyde-fixed; Triton X-100 at room temperature for 20 min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Antibody incubation with (IL-2R gamma) polyclonal Antibody, Unconjugated (bs-2545R) 1:100, 90 minutes at 37°C; followed by a conjugated Goat Anti-Rabbit IgG antibody at 37°C for 90 minutes, DAPI (blue, C02-04002) was used to stain the cell nuclei.



SHSY5Y cell; 4% Paraformaldehyde-fixed; Triton X-100 at room temperature for 20 min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Antibody incubation with (IL-2R gamma) polyclonal Antibody, Unconjugated (bs-2545R) 1:100, 90 minutes at 37°C; followed by a conjugated Goat Anti-Rabbit IgG antibody at 37°C for 90 minutes, DAPI (blue, C02-04002) was used to stain the cell nuclei.



Blank control: U937(blue). Primary Antibody: Rabbit Anti-IL-2R,gamma antibody(bs-2545R), Dilution: 1μg in 100 μL 1X PBS containing 0.5% BSA; Isotype Control Antibody: Rabbit IgG (orange) ,used under the same conditions. Secondary Antibody: Goat anti-rabbit IgG-PE(white blue), Dilution: 1:200 in 1 X PBS containing 0.5% BSA. Protocol The cells were fixed with 2% paraformaldehyde (10 min). Primary antibody (bs-2545R, 1µg/1x10^6 cells) were incubated for 30 min on the ice, followed by 1 X PBS containing 0.5% BSA + 10% goat serum (15 min) to block non-specific protein-protein interactions. Then the Goat Antirabbit IgG/PE antibody was added into the blocking buffer mentioned above to react with the primary antibody at 1/200 dilution for 30 min on ice. Acquisition of 20,000 events was performed.



Black line: Positive blank control (U937); Negative blank control (MCF7) Green line: Primary Antibody (Rabbit Anti-2R gamma antibody (bs-2545R)) Orange line: Isotype Control Antibody (Rabbit IgG) . Blue line : Secondary Antibody (Goat anti-rabbit IgG-AF647) U937 (Positive) and MCF7 (Negative control) cells (black) were incubated in 5% BSA blocking buffer for 30 min at room temperature. Cells were then stained with 2R gamma Antibody(bs-2545R)at 1:50 dilution in blocking buffer and incubated for 30 min at room temperature, washed twice with 2% BSA in PBS, followed by secondary antibody(blue) incubation for 40 min at room temperature. Acquisitions of 20,000 events were performed. Cells stained with primary antibody (green), and isotype control (orange).

- SELECTED CITATIONS -

• [IF=4.718] Arumugam P et al. Expression of a Functional IL-2 Receptor in Vascular Smooth Muscle Cells. The Journal of Immunology,2018 ji1701151. WB; Human. doi:10.4049/jimmunol.1701151